

Policy	MM-057
Effective Date	09/01/2024
Reviewed/Revised Date	4/27/2026
Next Review Date	4/27/2027
Origination Date	07/10/2024
Originated Department	Clinical Operations

Vitamin D Testing

Audience
Providers, Members, Brokers, MHC

Purpose
<p>Medical policies provide general support for applying Mountain Health Co-Op member policy document coverage decisions and must reference the member-specific benefit plan document. The terms of the member-specific Policy document may differ from the standard benefit plan on which this medical policy is based. If there is a conflict between a member-specific policy document and the Mountain Health Co-Op medical policy, the member-specific policy document supersedes this medical policy. Any person(s) applying this medical policy must identify member eligibility, the member-specific policy document, and related policies or guidelines before applying this medical policy, including the existence of any state or federal guidance. Mountain Health Co-Op medical policies are designed for informational purposes only and are not an authorization, explanation of benefits, or contract. Receipt of benefits is subject to satisfaction of all terms and conditions of the member-specific policy document coverage. Mountain Health Co-Op reserves the sole discretionary right to modify all policies and guidelines at any time.</p>

Definition
<p>Vitamin D is a fat-soluble vitamin. Very few foods naturally contain vitamin D (fatty fish and eggs are the exception), so vitamin D is obtained primarily through fortified foods or supplements and dermal synthesis from exposure to sunlight. Vitamin D has two forms, ergocalciferol (vitamin D2) and cholecalciferol (vitamin D3), and several metabolites. Estimates of vitamin D requirements vary and depend in part upon sun exposure and the standards used to define a deficient state. It is more accurate to consider fat-soluble vitamin D as a steroid hormone, synthesized by the skin and metabolized by the kidney to an active hormone, calcitriol. Clinical disorders related to vitamin D may arise because of altered</p>

availability of the parent vitamin D, altered conversion of vitamin D to its predominant metabolites, altered organ responsiveness to hydroxylated metabolites and disturbances in the interactions of the vitamin D metabolites with PTH and calcitonin.

Policy/Procedure

Commercial Plans/CHIP

Mountain Health Co-Op limits coverage for vitamin D (25-OH vitamin D) testing to specific circumstances demonstrated to impact health outcomes of its members.

1. Circumstances in which vitamin D (25-hydroxy vitamin D testing, (CPT® 82306) is covered:

1.1 Non-pregnant individual age 18 – 64 years with any of the following:

- a) Celiac Disease
- b) Chronic kidney disease stage III or greater
- c) Cirrhosis/Hepatic failure
- d) Hypocalcemia (persistent)
- e) Hypercalcemia (persistent)
- f) Hypercalciuria (persistent)
- g) Hypervitaminosis D
- h) Malabsorption syndromes including:
 - i. Cystic fibrosis;
 - ii. Inflammatory Bowel Disease - Crohn's disease;
 - iii. Bariatric surgery;
 - iv. Radiation Enteritis.
- i) Obstructive jaundice
- j) Osteomalacia
- k) Osteosclerosis/petrosis
- l) Osteoporosis if:
 - i. T score on DEXA scan; or
 - ii. History of fragility fractures; or
 - iii. FRAX > 3% 10-year probability of hip fracture or 20% 10-year probability of other major osteoporotic fracture; or
 - iv. FRAX > 3% (any fracture) with T-score v. Initiating bisphosphonate therapy (Vit D level should be determined and managed as necessary before bisphosphonate is initiated).
- m) Rickets, acquired or congenital childhood
- n) Vitamin D deficiency on replacement therapy related to a condition listed above; to monitor the efficacy of treatment
- o) Nephrolithiasis
- p) Hyper/hypoparathyroidism
- q) Chronic use of certain medications:
 - i. Antiseizure medications;
 - ii. Glucocorticoids;
 - iii. AIDS/HIV medications;
 - iv. Antifungals, e.g. ketoconazole;

- v. Cholestyramine.
- r) Granuloma-forming disorders:
 - i. Sarcoidosis;
 - ii. Tuberculosis;
 - iii. Histoplasmosis;
 - iv. Coccidiomycosis;
 - v. Berylliosis.

2. Conditions for which 1, 25-OH vitamin D level (CPT® 82652) is covered:

- 2.1 Unexplained hypercalcemia (suspected granulomatous disease or lymphoma);
- 2.2 Unexplained hypercalciuria (suspected granulomatous disease or lymphoma);
- 2.3 Suspected genetic childhood rickets;
- 2.4 Suspected tumor-induced osteomalacia;
- 2.5 Nephrolithiasis or hypercalciuria.

Mountain Health Co-Op does NOT cover vitamin D testing for any other indication including screening in the general population as it is considered not medically necessary.

Mountain Health Co-Op does NOT cover vitamin D testing (82306 or 82652 in any combination) more frequently than twice in a rolling 12-month period as it is considered not medically necessary for any diagnosis other than chronic kidney disease (CKD) or intestinal malabsorption.

Mountain Health Co-Op does NOT cover vitamin D testing utilizing both CPT® 82306 and CPT® 82652 in combination as it is considered not medically necessary.

3. Clinical Rationale

3.1 There is a paucity of evidence evaluating the benefit and harm of testing for vitamin D. Peer-reviewed scientific literature primarily investigates the effects of vitamin D supplementation, not testing. The Washington State Health Care Authority Health Technology Assessment Program published a technology assessment on vitamin D Screening and Testing in 2012. It was determined that no definitive conclusions can be drawn about the effectiveness of vitamin D screening or testing since no trials have been conducted to directly assess the impact of screening or testing on health outcomes, patient behavior, or clinical decision making. However, for some populations and outcomes, an association between serum levels and health outcomes and/or a positive effect of supplementation on health outcomes has been demonstrated. Thus, vitamin D screening has potential utility for identifying individuals who could benefit from the preventive or disease modifying effects of supplementation in these clinical situations when certain conditions are present or suspected to be present.

3.2 Subsequently, LeBlanc et al. in 2015 conducted a systematic review for the U.S. Preventive Services Task Force (USPSTF) to assess the benefits and harms of vitamin D screening in asymptomatic adults. The authors found “No study evaluated clinical outcomes or harms in persons screened versus not screened for vitamin D deficiency”. Limited evidence in persons not known to have conditions associated with vitamin D deficiency demonstrated that treating this deficiency with vitamin D may be associated with decreased risk for death in institutionalized elderly adults

and a reduction in the average number of falls but not fractures. The authors concluded that future research is needed to reduce assay variability; determine appropriate thresholds for vitamin D deficiency; and clarify effects of screening, subsequent treatment, and the subpopulations most likely to benefit. The USPSTF noted an “I” designation for this testing as routine screening outside of specific conditions being present. Many professional societies/organizations have also provided guidance related to the proper role of vitamin D testing. These include the Endocrine Society Clinical Practice Guideline on Evaluation, Treatment, and Prevention of vitamin D Deficiency (Holick, et al., 2011) which recommends specific testing for vitamin D deficiency in individuals at risk for deficiency and do not recommend population screening for vitamin D deficiency in individuals who are not at risk. These guidelines also recommend using the serum circulating 25-hydroxyvitamin D [25(OH)D] level, measured by a reliable assay, to evaluate vitamin D status in patients who are at risk for vitamin D deficiency. As for serum 1,25-dihydroxyvitamin D [1,25(OH)2D] assay these guidelines do not recommend using this testing for screening (patients at risk) and are in favor of using it only in monitoring certain conditions, such as acquired and inherited disorders of vitamin D and phosphate metabolism. The authors note there is no evidence demonstrating benefits of screening for vitamin D deficiency at a population level. Such evidence would require demonstration of the feasibility and cost-effectiveness of such a screening strategy, as well as benefits in terms of important health outcomes. In the absence of this evidence, it is premature to recommend screening at large at this time. Currently, 25(OH)D measurement is reasonable to use in groups of people at high risk for vitamin D deficiency and in whom a prompt response to optimization of vitamin D status could be expected.

- 3.3** The American Academy of Pediatrics (AAP) Committee on Nutrition (Golden, et al., 2014) states that evidence is insufficient to recommend universal screening for vitamin D deficiency. The AAP report advises screening for vitamin D deficiency “only in children and adolescents with conditions associated with reduced bone mass and/or recurrent low-impact fractures. More evidence is needed before recommendations can be made regarding screening of healthy black and Hispanic children or children with obesity. The recommended screening is measuring serum 25-OH-D concentration, and it is important to be sure this test is chosen instead of measurement of the 1,25-OH2-D concentration, which has little, if any, predictive value related to bone health.”
- 3.4** Whereas the American Association of Clinical Endocrinologists (AACE) and American College of Endocrinology (ACE) have published a Clinical Practice Guidelines for the Diagnosis and Treatment of Postmenopausal Osteoporosis (Camacho, et al., 2016) which includes the following recommendation for fundamental measures concerning bone health: Measure serum 25-hydroxyvitamin D (25[OH]D) in patients who are at risk for vitamin D insufficiency, particularly those with osteoporosis (Grade B).
- 3.5** Several societies including the AACE/ACE, The Obesity Society, American Society for Metabolic & Bariatric Surgery (ASMBS), Obesity Medicine Association, and American Society of Anesthesiologists in 2020 provided evidence-based recommendations for Perioperative nutrition, metabolic, and nonsurgical support of

patients undergoing Bariatric procedures. They recommend baseline and annual postoperative evaluation for vitamin D deficiency is recommended after Roux-en-Y gastric bypass, sleeve gastrectomy, or laparoscopic biliopancreatic diversion without or with duodenal switch (Recommendation 53) (Mechanick, 2019). The American Society for Metabolic and Bariatric Surgery (ASMBS) Integrated Health Nutritional Guidelines for the Surgical Weight Loss Patient 2016 Update: Micronutrients (Parrott, et al., 2017) specifically recommends routine pre- and post-weight loss surgery (WLS) vitamin D screening. The ASMBS states routine post-WLS screening refers to performing a nutrient assessment every 3–6 months in the first year and annually thereafter, unless otherwise specified.

- 3.6** As for circumstances involving primary hyperparathyroidism, Wilhelm, et al. in 2016 writing for the American Association of Endocrine Surgeons (AAES): made the recommendation to perform biochemical evaluation of suspected primary hyperparathyroidism including serum total calcium, PTH, creatinine, and 25-hydroxyvitamin D levels (strong recommendation; moderate quality evidence).
- 3.7** In cases of Crohn’s disease the American College of Gastroenterology (ACG) recommends including “Routine laboratory investigation: Initial laboratory investigation should include evaluation for inflammation, anemia, dehydration, and malnutrition” (Lichtenstein, et al., 2018). This includes testing for 25(OH) vitamin D. The ACG also make recommendations in assessment of Primary Sclerosing Cholangitis (Lindor, et al., 2015) to screen and monitor patients with advanced liver disease should be for fat-soluble vitamin deficiencies as fat-soluble vitamin deficiencies can occur in late stages of primary sclerosing cholangitis when patient becomes jaundiced. Levels of vitamins A, E, and D should be assessed in patients with advanced disease (Conditional recommendation, moderate quality of evidence).
- 3.8** The ACG Clinical Guideline on the Diagnosis and Management of Celiac Disease (Rubio-Tapia, et al., 2013) recommends people with newly diagnosed celiac disease should undergo testing and treatment for micronutrient deficiencies. Deficiencies to be considered for testing should include, but not be limited to, iron, folic acid, vitamin D, and vitamin B12. (Conditional recommendation, low level of evidence).
- 3.9** Evidence-based guidelines from the National Osteoporosis Foundation (NOF): The NOF’s Clinician’s Guide to Prevention and Treatment of Osteoporosis (Cosman, et al., 2014) notes serum 25- hydroxyvitamin D (25[OH]D) testing should be performed to evaluate for secondary causes of osteoporosis. It also states, “since vitamin D intakes required to correct vitamin D deficiency are so variable among individuals, serum 25(OH)D levels should be measured in patients at risk of deficiency. Vitamin D supplements should be recommended in amounts sufficient to bring the serum 25(OH)D level to approximately 30 ng/ml (75 nmol/L) and a maintenance dose recommended to maintain this level, particularly for individuals with osteoporosis. It defines vitamin D insufficiency as a serum 25- hydroxyvitamin D (25[OH]D)<30 ng/ml (75 nmol/L).

Applicable Coding

CPT Codes

Covered Codes

82306 Vitamin D; 25 hydroxy, includes fraction(s), if performed 82652 Vitamin D; 1, 25 dihydroxy, includes fraction(s), if performed

Not Covered Codes

0038U Vitamin D, 25 hydroxy D2 and D3, by LC-MS/MS, serum microsample, quantitative

HCPCS Codes

No applicable HCPCS codes

ICD-10 Codes

A15.0-A19.9 Tuberculosis

C22.0-C22.9 Malignant neoplasm of liver and intrahepatic bile ducts

C23 Malignant neoplasm of gallbladder

C24.0-C24.9 Malignant neoplasm of other and unspecified parts of biliary tract

C25.0-C25.9 Malignant neoplasm of pancreas

C26.0-C26.9 Malignant neoplasm of other and ill-defined digestive organs

C83.80-C83.89 Other non-follicular lymphoma

C84.00-C84.09 Mycosis fungoides

C84.10-C84.19 Sezary disease

C88.4 Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue [MALT-lymphoma]

D13.0-D13.9 Benign neoplasm of other and illdefined parts of digestive system

D86.0-D86.89 Sarcoidosis

E20.0 Idiopathic hypoparathyroidism

E20.8 Other hypoparathyroidism

E20.9 Hypoparathyroidism, unspecified

E21.0-E21.5 Hyperparathyroidism and other disorders of parathyroid gland

E24.0-E24.9 Cushing's syndrome E41 Nutritional marasmus E55.0-E55.9 Vitamin D deficiency

E64.3 Sequelae of rickets

E67.2 Megavitamin-B6 syndrome

E67.3 Hypervitaminosis D

E67.8 Other specified hyperalimentation

E68 Sequelae of hyperalimentation

E83.30-E83.39 Disorders of phosphorus metabolism and phosphatases

E83.50-E83.59 Disorders of calcium metabolism

E84.0-E84.9 Cystic fibrosis

E89.2 Postprocedural hypoparathyroidism K50.00-K50.919 Crohn's disease [regional enteritis]

K70.2 Alcoholic fibrosis and sclerosis of liver K70.30-K70.31 Alcoholic cirrhosis of liver

K74.0-K74.69 Fibrosis and cirrhosis of liver

K75.81 Nonalcoholic steatohepatitis (NASH)

K76.0 Fatty (change of) liver, not elsewhere classified

K76.89 Other specified diseases of live

K82.0 Obstruction of gallbladder

K82.8 Other specified diseases of gallbladder K82.9 Disease of gallbladder, unspecified

K83.01-K83.9 Other diseases of biliary tract

K85.10-K85.12 Biliary acute pancreatitis

K86.2 Cyst of pancreas

K86.3 Pseudocyst of pancreas

K86.81-K86.89 Other specified diseases of pancreas

K86.9 Disease of pancreas, unspecified
K87 Disorders of gallbladder, biliary tract and pancreas in diseases classified elsewhere
K90.0 Celiac Disease
K90.0-K90.49 Intestinal malabsorption
K90.89 Other intestinal malabsorption
K90.9 Intestinal malabsorption, unspecified
K91.2 Postsurgical malabsorption, not elsewhere classified
L90.0 Lichen sclerosus et atrophicus
L94.0 Localized scleroderma [morphea]
L94.1 Linear scleroderma
L94.3 Sclerodactyl
M05.00-M05.9 Rheumatoid arthritis with rheumatoid factor
M06.00-M06.9 Other rheumatoid arthritis
M32.0-M32.9 Systemic lupus erythematosus (SLE)
M33.01-M33.09 Juvenile dermatomyositis M33.11-M33.19 Other dermatomyositis M33.91-
M33.99 Dermatopolymyositis
M36.0 Dermato(poly)myositis in neoplastic disease M81.0-M81.8 Osteoporosis without
current pathological fracture M83.0-M83.9 Adult osteomalacia
M85.80-M85.9 Other specified disorders of bone density and structure, various sites
M88.0-M88.9 Osteitis deformans of various sites
N18.3 Chronic kidney disease (CKD), stage III N18.4 Chronic kidney disease (CKD), stage IV
N18.5 Chronic kidney disease (CKD), Stage V
N25.81 Secondary hyperparathyroidism of renal origin
Q78.0 Osteogenesis imperfecta
Q78.2 Osteopetrosis

Vendors

- **Personify**
- **HPS**

References

1. ACOG Committee on Obstetric Practice. ACOG Committee Opinion No. 495: Vitamin D: Screening and supplementation during pregnancy. Obstet Gynecol. 2011 Jul;118(1):197-8. Reaffirmed 2019. Accessed June 2020. Available at URL address: <https://www.acog.org>
2. American Academy of Neurology. Policy and Guidelines. Accessed Jan 2020. Available at URL address: <https://www.aan.com/policy-and-guidelines/> and <https://www.aan.com/Guidelines/Home/ByStatusOrType?status=all>
3. American Association of Clinical Endocrinologists. AACE/ACE CLINICAL PRACTICE GUIDELINES. Accessed June 2020. Available at URL address: <https://www.aace.com>
4. American Association of Endocrine Surgeons. Practice Guidelines. Accessed June 2020. Available at URL address: <https://www.endocrinesurgery.org>
5. American Association of Endocrine Surgeons. Practice Guidelines. Accessed June 2020. Available at URL address: <https://www.endocrinesurgery.org>
6. American College of Rheumatology. Clinical Practice Guidelines. Accessed June 2020. Available at URL address: <https://www.rheumatology.org/Practice-Quality/Clinical-Support/Clinical-Practice-Guidelines>
7. American Gastroenterological Association. Accessed June 2020. Available at URL address: <https://www.gastro.org/guidelines>

8. American Geriatrics Society Workgroup on Vitamin D Supplementation for Older Adults. Recommendations abstracted from the American Geriatrics Society Consensus Statement on vitamin D for Prevention of Falls and Their Consequences. *J Am Geriatr Soc.* 2014 Jan;62(1):147-52. Accessed Jan 2020. Status: INACTIVE. Available at URL address: <https://geriatricscareonline.org/ProductAbstract/american-geriatrics-society-consensus-statement-vitamin-d-for-prevention-of-falls-and-their-consequences-in-older-adults/CL009/?param2=search>
9. American Thyroid Association. Guidelines and Surgical Statements. Accessed June 2020. Available at URL address: <https://www.thyroid.org/professionals/ata-professional-guidelines/>
10. Camacho PM, Petak SM, Binkley N, Clarke BL, Harris ST, Hurley DL, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of Postmenopausal Osteoporosis - 2016--executive summary. *Endocr Pract.* 2016 Sep;22(9):1111-8.
11. Centers for Medicare & Medicaid Services (CMS). Local Coverage Determinations (LCDs) Alphabetical Index. Vitamin D Assay Testing. Accessed June 2020. Available at URL address: <https://med.noridianmedicare.com>
12. Charoenngam, N. (2021). "Vitamin D and Rheumatic Diseases: A Review of Clinical Evidence." *Int J Mol Sci* 22(19).
13. Choosing Wisely® Initiative. Accessed Jan 2020. Available at URL address: <http://www.choosingwisely.org/clinician-lists/>
14. Cosentino, N., et al. (2021). "Vitamin D and Cardiovascular Disease: Current Evidence and Future Perspectives." *Nutrients* 13(10).
15. Cosman, F., et al. (2014). "Clinician's Guide to Prevention and Treatment of Osteoporosis." *Osteoporos Int* 25(10): 2359- 2381. Accessed June 2020.
16. Golden, N. H., et al. (2014). "Optimizing bone health in children and adolescents." *Pediatrics* 134(4): e1229-1243.
17. Holick, M. F., et al. (2011). "Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline." *J Clin Endocrinol Metab* 96(7): 1911-1930.
18. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, American College of Cardiology/American Heart Association Task Force on Practice Guidelines; Obesity Society, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol.* 2014 Jul 1;63(25 Pt B):2985-3023. Erratum in *J Am Coll Cardiol.* 2014 Jul 1;63(25 Pt B):3029-3030.
19. Kahwati LC, LeBlanc E, Weber RP, Giger K, Clark R, Suvada K, Guisinger A, Viswanathan M. Screening for Vitamin D Deficiency in Adults: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA.* 2021 Apr 13;325(14):1443-1463. doi: 10.1001/jama.2020.26498. PMID: 33847712.
20. LeBlanc, E. S., et al. (2015). "Screening for vitamin D deficiency: a systematic review for the U.S. Preventive Services Task Force." *Ann Intern Med* 162(2): 109-122.
21. Lichtenstein, G. R., et al. (2018). "ACG Clinical Guideline: Management of Crohn's Disease in Adults." *Am J Gastroenterol* 113(4): 481-517.
22. Lindor, K. D., et al. (2015). "ACG Clinical Guideline: Primary Sclerosing Cholangitis." *Am J Gastroenterol* 110(5): 646-659; quiz 660.
23. Mechanick, J. I., et al. (2019). "Clinical Practice Guidelines for the Perioperative Nutrition, Metabolic, and Nonsurgical Support of Patients Undergoing Bariatric Procedures - 2019 Update: Cosponsored by American Association of Clinical Endocrinologists/American College of Endocrinology, the Obesity Society, American Society for Metabolic & Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists - Executive Summary." *Endocr Pract* 25(12): 1346-1359.
24. Parrott, J., et al. (2017). "American Society for Metabolic and Bariatric Surgery Integrated Health Nutritional Guidelines for the Surgical Weight Loss Patient 2016 Update: Micronutrients." *Surg Obes Relat Dis* 13(5): 727-741.
25. Rubio-Tapia, A., et al. (2013). "ACG clinical guidelines: diagnosis and management of celiac disease." *Am J Gastroenterol* 108(5): 656-676; quiz 677.
26. Singer AG, McChesney C. Reduce unnecessary routine vitamin D testing. *Can Fam Physician.* 2023 Sep;69(9):620-622. doi: 10.46747/cfp.6909620. PMID: 37704241; PMCID: PMC10498906.

27. UpToDate® (2023). "Management of celiac disease in adults". Literature current through April 2023. Topic last updated: February 1, 2023. Version 43.0. Topic 4777. Accessed: May 31, 2023. Available at: <https://www.uptodate.com>
28. UpToDate® (2023). "Vitamin D deficiency in adults: Definition, clinical manifestations, and treatment". Literature current through March 2024. Topic last updated: September 30, 2023. Version 69.0. Topic 2022. Accessed: April 16, 2024. Available at: <https://www.uptodate.com>
29. UpToDate® (2022). "Vitamin D insufficiency and deficiency in children and adolescents". Literature current through April 2023. Topic last updated: April 12, 2022. Version 59.0. Topic 14590. Accessed: May 31, 2023. Available at: <https://www.uptodate.com>
30. U.S. Preventive Services Task Force (USPSTF). Final Recommendation Statement Vitamin D Deficiency: Screening. November 2021. Accessed June 12, 2022. Available at URL address: <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/vitamin-d-deficiency-screening>
31. U.S. Preventive Services Task Force (USPSTF). Final Recommendation Statement Vitamin D, Calcium, or Combined Supplementation for the Primary Prevention of Fractures in Community-Dwelling Adults: Preventive Medication. April 2018. Accessed June 2020. Available at URL address: <https://www.uspreventiveservicestaskforce.org>
32. Washington State Health Care Authority. Health Technology Assessment Program (HTA). Vitamin D Screening and Testing. Final Evidence Report November 16, 2012. Accessed June 2020. Available at URL address: [https://www.hca.wa.gov/assets/program/vitd_finalrpt_111612\[1\].pdf](https://www.hca.wa.gov/assets/program/vitd_finalrpt_111612[1].pdf)
33. Wilhelm, S. M., et al. (2016). "The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism." JAMA Surg 151(10): 959-968.

Review/Revision/Approval History	
Date	Description
07/10/2024	New Policy
04/27/2026	Reviewed and approved by Mountain Health CO-OP Policy Committee

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult appropriate healthcare providers for medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member’s benefit plan, effective when services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Including or excluding a procedure, diagnosis, or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as they apply to an individual member.

Mountain Health Co-Op makes no representations and accepts no liability regarding the content of any external information cited or relied upon in this policy. Mountain Health Co-Op updates its Coverage Policies regularly and reserves the right to amend these policies and give notice per State and Federal requirements. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Mountain Health Co-Op.

”Mountain Health Co-Op” and its accompanying logo and marks are protected and registered trademarks of Mountain Health Co-Op. The content of this Service is proprietary and protected by copyright. You may access the copyrighted content of this Service only for purposes outlined in these Conditions of Use.

© CPT Only – American Medical Association