

Policy	MM-086
Effective Date	09/01/2024
Reviewed/Revised Date	04/27/2026
Next Review Date	04/27/2027
Origination Date	07/01/2024
Originated Department	Clinical Operations

Automated Percutaneous and Percutaneous Endoscopic Discectomy

Audience
Medical Management

Purpose
<p>Medical policies provide general support for applying Mountain Health Co-Op member policy document coverage decisions and must reference the member-specific benefit plan document. The terms of the member-specific Policy document may differ from the standard benefit plan on which this medical policy is based. If there is a conflict between a member-specific policy document and the Mountain Health Co-Op medical policy, the member-specific policy document supersedes this medical policy. Any person(s) applying this medical policy must identify member eligibility, the member-specific policy document, and related policies or guidelines before applying this medical policy, including the existence of any state or federal guidance. Mountain Health Co-Op medical policies are designed for informational purposes only and are not an authorization, explanation of benefits, or contract. Receipt of benefits is subject to satisfaction of all terms and conditions of the member-specific policy document coverage. Mountain Health Co-Op reserves the sole discretionary right to modify all policies and guidelines at any time.</p>

Definition
N/A

Policy/Procedure
<p>Policy Statement and Criteria</p> <p>Commercial Plans</p> <p>Mountain health Co-Op considers services and procedures listed in the current and future automated percutaneous discectomy is considered investigational as a technique of</p>

intervertebral disc decompression in individuals with back pain and/or radiculopathy related to disc herniation in the lumbar, thoracic, or cervical spine.

Percutaneous endoscopic discectomy is considered investigational as a technique of intervertebral disc decompression in individuals with back pain and/or radiculopathy related to disc herniation in the lumbar, thoracic, or cervical spine

Section 1862(a)(1)(A) of the Social Security Act is the basis for denying payment for types of care, specific items, services, or procedures, not excluded by any other statutory clause, meeting all technical requirements for coverage, but are determined to be any of the following:

- Not generally accepted in the medical community as safe and effective in the setting and for the condition for which it is used,
- Not proven to be safe and effective based on peer review or scientific literature
- Experimental
- Not medically necessary in the particular case
- Furnished at a level, duration or frequency that is not medically appropriate
- Not furnished in accordance with accepted standards of medical practice, or
- Not furnished in a setting appropriate to the patient's medical needs and condition.

Items and services must be established as safe and effective to be considered medically necessary. That is, the items and services must be:

- Consistent with the symptoms or diagnosis of the illness or injury under treatment; and
- Necessary for, and consistent with, generally accepted professional medical standards of care (e.g., not experimental or investigational);and
- Not furnished primarily for the convenience of the patient, the provider or supplier; and
- Furnished at the most appropriate level that can be provided safely and effectively to the patient.

Medical devices that are not approved for marketing by the Food and Drug Administration (FDA) are considered investigational by Medicare and are not considered reasonable and necessary for the diagnosis or treatment of illness or injury, or to improve functioning of a malformed body member. Mountain Health Co-Op payments, therefore, may not be made for medical procedures and services performed using devices that have not been approved for marketing by the FDA or for those not included in an FDA-approved investigational (IDE) trial.

Background

Back pain or radiculopathy related to herniated discs is an extremely common condition and a frequent cause of chronic disability. Although many cases of acute low back pain and radiculopathy will resolve with conservative care, surgical decompression is often considered when the pain is unimproved after several months and is clearly neuropathic in origin, resulting from irritation of the nerve roots. Open surgical treatment typically consists of discectomy in which the extruding disc material is excised. When performed with an operating microscope, the procedure is known as a microdiscectomy.

Minimally invasive options have also been researched, in which some portion of the disc is removed or ablated, although these techniques are not precisely targeted at the offending extruding disc material. Ablative techniques include laser discectomy and radiofrequency decompression (see evidence review 7.01.93). Intradiscal electrothermal annuloplasty is another minimally invasive approach to low back pain. In this technique, radiofrequency energy is used to treat the surrounding disc annulus (see evidence review 7.01.72).

Herein, BCBSA addresses automated percutaneous and endoscopic discectomy, in which the disc decompression is accomplished by the physical removal of disc material rather than its ablation. Traditionally, discectomy was performed manually through an open incision, using cutting forceps to remove nuclear material from within the disc annulus. This technique was modified by automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Endoscopic techniques may be intradiscal or may involve extraction of noncontained and sequestered disc fragments from inside the spinal canal using an interlaminar or transforaminal approach. Following insertion of the endoscope, decompression is performed under visual control.

Regulatory Status

The Dekompressor[®] Percutaneous Discectomy Probe (Stryker), Herniatome Percutaneous Discectomy Device (Gallini Medical Devices), and the Nucleotome[®] (Clarus Medical) are examples of percutaneous discectomy devices that have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA indication for these products is for “aspiration of disc material during percutaneous discectomies in the lumbar, thoracic and cervical regions of the spine.” FDA product code: HRX.

A variety of endoscopes and associated surgical instruments have also been cleared for marketing by FDA through the 510(k) process.

Rationale

This evidence review was created in December 1995 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through May 5, 2023.

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific

outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

Automated Percutaneous Discectomy

Clinical Context and Therapy Purpose

The purpose of automated percutaneous discectomy is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with herniated intervertebral disc(s).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with herniated intervertebral disc(s).

Interventions

The therapy being considered is automated percutaneous discectomy.

Comparators

The following therapies and practices are currently being used to treat herniated intervertebral disc(s): conservative therapy and open discectomy or microdiscectomy.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific outcomes measured by specific instruments may include improvements in functional outcomes assessed on the Oswestry Disability Index (ODI), reductions in pain using a visual analog scale (VAS), improvements in quality of life measured on the 36-Item Short-Form Health Survey (SF-36) and Euro-QOL-5D, and treatment-related morbidity including surgical success/failure and complications. To assess outcomes, follow-up at 1 year is considered appropriate.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

Systematic reviews have assessed automated percutaneous discectomy compared to other interventions; however, the majority of these reviews contained observational studies published more than a decade ago with generally small patient populations and inconsistent results. Lewis et al (2015) published the most recent systematic review and network meta-analysis comparing trials of 21 different treatment strategies for sciatica.¹ Examples of the 21 treatment strategies included in the analysis include conservative care, disc surgery, intraoperative interventions, epidural injections, biologic agents, and percutaneous discectomy. Under the category of “percutaneous discectomy,” reviewers combined automated percutaneous discectomy, percutaneous automated nucleotomy, nucleoplasty, and laser discectomy. They searched 28 databases and trial registries through December 2009. Ninety studies were included and 10 involved the percutaneous discectomy category as an intervention. Of the 10, 4 are relevant to this evidence review: 2 case-control studies of percutaneous endoscopic discectomy (2006, 2007), 1 RCT of percutaneous endoscopic

discectomy (1993), and 1 RCT of automated percutaneous discectomy (1995). The remaining studies were published in a foreign language or involved other comparators (nucleolysis, chemonucleolysis). The global effects odds ratio for the category of percutaneous discectomy compared with inactive control was 0.82 (95% confidence interval [CI], 0.39 to 1.72), which was inferior to disc surgery, epidural injections, and intraoperative interventions. The pain intensity weighted mean difference for the category of percutaneous discectomy compared with inactive control was 11.5 (95%, -18.6 to 41.6). Reviewers concluded that there was no support for the effectiveness of percutaneous discectomy for the treatment of sciatica. Due to the inclusion of additional interventions into the broad category of percutaneous discectomy in this review, the relevance of these results to this evidence review is limited.

Randomized Controlled Trials

The 2002 Lumbar Automated Percutaneous Discectomy Outcomes Group (LAPDOG) trial is a RCT to compare automated percutaneous discectomy with open discectomy in patients with lumbar disc herniation.² No additional RCTs have been identified since the 2002 LAPDOG trial. The trial was designed to recruit 330 patients but enrolled 36 patients for reasons not readily apparent. Twenty-seven patients were available at follow-up, with efficacy reported by 41% of those undergoing automated percutaneous discectomy and by 40% of those undergoing conventional discectomy. The trialists concluded that “It is difficult to understand the remarkable persistence of percutaneous discectomy in the face of a virtually complete lack of scientific support for its effectiveness in treated lumbar disc herniation.” The tables below more fully describe key characteristics, results, and limitations of the LAPDOG trial.

Table 1. Characteristics of the LAPDOG Trial

Study	Countries	Sites	Dates	Participants	Interventions
Haines et al (2002) ²	US, Canada	10	NR	Patients with predominantly unilateral leg pain or paresthesia with no previous treatment for lumbar spinal disease, at least 2 of 4 objective signs, and an imaging study confirming disc herniation at the appropriate level	Automated percutaneous discectomy vs. conventional discectomy

OG: Lumbar Automated Percutaneous Discectomy Outcomes Group; NR: not reported.

Table 2. Results of the LAPDOG Trial

Study	Treatment success ^a (at 6 months)	Treatment failure ^b (at 6 months)	SF-36 Physical Functioning Subscore	SF-36 General Health Subscore	Modified Roland Score
Haines et al (2002) ²					
N	27	27	NR	NR	NR
Automated percutaneous discectomy,	7 (41%)	10 (59%)	Pre- vs. postoperative mean difference: 35.7	Pre- vs. postoperative mean difference: 5.0	Pre- vs. postoperative mean difference: 9.7
Conventional discectomy	4 (40%)	6 (60%)	Pre- vs. postoperative mean difference: 36.1	Pre- vs. postoperative mean difference: 8.0	Pre- vs. postoperative mean difference: 10.6
p	.95	.95	.96	.58	.74

LAPDOG: Lumbar Automated Percutaneous Discectomy Outcomes Group; NR: not reported; SF-36: 36-Item Short-Form Health Survey.

^aSuccess was defined as either an excellent or good result as defined by an outcome matrix.

^bFailure was defined as not achieving success or requiring a second procedure during the follow-up period.

Table 3. Study Relevance Limitations of the LAPDOG Trial

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Haines et al (2002) ²	4-3. Investigators believed that study inclusion criteria reflected an existing population with lumbar disc			4. Primary outcomes of “success” or “failure” largely subjective in nature; investigators admit that the	1,2. Outcomes reported only for 6 months of follow-up; 12 month follow-up was achieved for only 19 patients and the study

	disease; however, results from only 27 patients were eventually analyzed from a planned enrollment of 330 patients			Outcome measurement tool used can not be precisely reproduced	did not report any of those results
--	--	--	--	---	-------------------------------------

LAPDOG: Lumbar Automated Percutaneous Discectomy Outcomes Group. The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other. ^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other. ^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other. ^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 4. Study Design and Conduct Limitations of the LAPDOG Trial

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Haines et al (2002) ²		1,2. Blinding did not appear to occur		1. Of 34 initially randomized patients, 9 were lost to follow-up, 6	3. Power estimates led the investigators to plan enrollment of	1. Beyond the cursory discussion of lack of power, a discussion of
				month follow-up data was obtained on only 27 patients, and 12 month follow-up data was obtained for only 19 patients	330 patients in order to reliably identify a difference in success rate of 15% or greater; results were analyzed on 27 patients	the statistical analyses is nonexistent

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. ^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias. ^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. ^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). ^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference. ^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for t; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

All published trials have focused on lumbar disc herniation. There were no RCTs of automated percutaneous discectomy for cervical or thoracic disc herniation. A review of the evidence from American Society of Interventional Pain Physicians (2013) noted that “even though Dekompessor [disc removal system] may be considered a new interventional modality, the early studies were published approximately 8 years ago. Consequently, one would expect that the technique’s continued use would be supported by more recent, high-quality evaluations.”³,

Section Summary: Automated Percutaneous Discectomy

The evidence for automated percutaneous discectomy in individuals who have herniated intervertebral disc(s) includes small RCTs and systematic reviews. Evidence from small RCTs does not support the use of this procedure. Well-designed and executed RCTs are needed to determine the benefits and risks of this procedure.

Percutaneous Endoscopic Discectomy

Clinical Context and Therapy Purpose

The purpose of percutaneous endoscopic discectomy is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with herniated intervertebral disc(s).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with herniated intervertebral disc(s).

Interventions

The therapy being considered is percutaneous endoscopic discectomy.

Comparators

The following therapies and practices are currently being used to treat herniated intervertebral disc(s): conservative therapy and open discectomy or microdiscectomy.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific outcomes measured by specific instruments include improvements in functional outcomes assessed on the ODI, reductions in pain using a VAS, improvements in quality of life measured on the SF-36 and Euro-QOL-5D, and treatment-related morbidity including surgical success/failure and complications. To assess outcomes, follow-up at 1 year is considered appropriate.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

A number of systematic reviews have evaluated the efficacy and safety of percutaneous endoscopic discectomy compared to open discectomy or microendoscopic discectomy. A comparison of the trials included in more recent systematic reviews (2017 to present) is shown in Table 5. Characteristics and results of these reviews are summarized in Tables 6 and 7.

Table 5. Trials Included in Systematic Reviews of Percutaneous Endoscopic Discectomy Versus Other Discectomy Procedures

Trials	Systematic Reviews							
	Phan et al (2017) ⁴ ,	Shi et al (2019) ⁵ ,	Yu et al (2019) ⁶ ,	Zhou et al (2020) ⁷ ,	Xu et al (2020) ⁸ ,	Bai et al (2021) ⁹ ,	Gadjradj et al (2021) ¹⁰ ,	Zhao et al (2022) ¹¹ ,
Ma et al (2022) ¹² ,								●
Wang et al (2021) ¹³ ,								●
Rajamani et al (2021) ¹⁴ ,								●
Jing et al (2021) ¹⁵ ,								●
Jarebi et al (2021) ¹⁶ ,								●
Meyer et al (2020) ¹⁷ ,								●
Chen et al (2020) ¹⁸ ,								●
Kim et al (2019) ¹⁹ ,								●

Ahn et al (2019) ²⁰ ,								●
Liu et al (2018) ²¹ ,								●
Sun et al (2017) ²² ,								●
Jeong et al (2006) ²³ ,								●
Akcakaya et al (2016) ²⁴ ,							●	
Choi et al (2018) ²⁵ ,							●	
Dai et al (2020) ²⁶ ,							●	
Krappel et al (2001) ²⁷ ,							●	
Tacconi et al (2019) ²⁸ ,							●	
Tacconi et al (2020) ²⁹							●	
Tao et al (2018) ³⁰							●	
Wang et al (2017) ³¹							●	
Xu et al (2017) ³²							●	
Ahn et al (2016) ³³						●		●
Chang et al (2018) ³⁴						●	●	●
Lui et al (2017) ³⁵						●		●
Pan et al (2016) ³⁶						●	●	●
Yao et al (2017) ³⁷						●		
Yao et al (2017) ³⁸						●		●
Gibson et al (2017) ³⁹				●			●	●
Hsu et al (2013) ⁴⁰				●				●
Kim et al (2007) ⁴¹				●		●		●
Qu et al (2017) ⁴²				●				
Wang et al (2013) ⁴³				●				●
Zhao et al (2012) ⁴⁴				●				

Yoon et al (2012) ⁴⁵	●	●	●		●			
Li et al (2015) ⁴⁶	●				●			●
Sinkemani et al (2015) ⁴⁷	●	●	●		●			●
Song et al (2017) ⁴⁸		●	●		●			●
Tu et al (2017) ⁴⁹					●			
Liu et al (2018) ²¹		●	●	●	●	●		
Li et al (2018) ⁵⁰		●	●	●	●			●
Abdurexiti et al (2018) ⁵¹		●	●		●			
Chen et al (2018) ⁵²		●	●	●	●	●		●
Lui et al (2012) ⁵³			●					
Wu et al (2009) ⁵⁴		●						
Yang et al (2015) ⁵⁵		●		●				
Duan et al (2016) ⁵⁶		●						
Zhao et al (2016) ⁵⁷		●						
Ding et al (2017) ⁵⁸		●						
Li et al (2017) ⁵⁹		●						
Lui et al (2017) ⁶⁰		●						
Luo et al (2017) ⁶¹		●						
Qu et al (2017) ⁶²		●						
Chen et al (2018) ⁶³		●						
Wu et al (2018) ⁶⁴		●						
Belykh et al (2016) ⁶⁵		●						
Chen et al (2015) ⁶⁶	●							●
Choi et al (2016) ⁶⁷	●			●				●
Garg et al (2011) ⁶⁸	●							

Hermantin et al (1999) ⁶⁹	●					●	
Huang et al (2005) ⁷⁰	●						
Hussein et al (2014) ⁷¹	●						
Kleinpeter et al (1995) ⁷²	●						
Lee et al (2009) ⁷³	●				●		●
Martin Laez et al (2012) ⁷⁴	●						
Mayer et al (1993) ⁷⁵	●		●		●	●	●
Ohya et al (2016) ⁷⁶	●						
Pan et al (2014) ⁷⁷	●						●
Righesso et al (2007) ⁷⁸	●						
Ruetten et al (2008) ⁷⁹	●						
Ruetten et al (2009) ⁸⁰	●				●		
Sasaoka et al (2006) ⁸¹	●						
Schizas et al (2005) ⁸²	●						
Teli et al (2010) ⁸³	●						
Ruetten et al (2007) ⁸⁴	●						
Ruetten et al (2008) ⁸⁵					●		
Lee et al (2006) ⁸⁶					●		

Table 6. Summary of Systematic Reviews of Percutaneous Endoscopic Discectomy Versus Other Discectomy Procedures

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Zhao et al (2022) ¹¹ ,	To May 2022	33	Patients with lumbar disc herniation who underwent PTED, MED or other surgical procedures	6467 (20-1856)	7 RCTs; 26 non-randomized controlled retrospective studies	Not reported

Bai et al (2021)⁹	To February 2018	14	Patients with lumbar disc herniation who underwent PELD or other surgical procedures	2528 (74-902)	4 RCTs; 10 cohort studies	Not reported
Gadjradj et al (2021)¹⁰	To April 2020	14	Patients with lumbar disc herniation who underwent PTED or open microdiscectomy	1465 (30-462)	9 RCTs; 5 prospective nonrandomized comparative studies	Follow-up: 3 to 12 months
Xu et al (2020)⁸	Search dates not stated; included trials from 2012 to 2018	9	Patients with single-level lumbar disc herniation who underwent PELD or MED for treatment	984 (51-216)	1 RCT ; 8 retrospective nonrandomized comparative studies	Follow-up: 1 to > 6 years
Zhou et al (2020)⁷	To October 2018	12	Patients with lumbar disc herniation who underwent PELD or MED for treatment	2400 (40-915)	4 RCTs; 8 retrospective nonrandomized comparative studies	Follow-up: 3 to 46 months
Yu et al (2019)⁶	To August 31, 2018	8	Patients with lumbar disc herniation who underwent PTED or MED procedures and were followed for at least 6 months	805 (51-216)	1 RCT ; 7 observational studies	Follow-up: 6 months to 5 years
			who underwent PELD or MED for treatment			
Phan et al (2017)⁴	To February 2016	23	Patients who underwent either an endoscopic or open approach for disc herniation; the endoscopic approach consisted of patients who underwent either FED or MED while the open approach included those who underwent open discectomy or microdiscectomy	28,487 (20-26,612)	10 RCTs; 4 prospective observational studies; 9 retrospective observational studies	Follow-up: 3 to 104 months

FED: full-endoscopic technique discectomy; MED: microendoscopic discectomy; PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trial

Table 7. Results of Systematic Reviews of Trials of Percutaneous Endoscopic Discectomy Versus Other Discectomy Procedures

Study	Length of stay	Leg pain VAS	Lower back pain VAS	ODI	Overall complication rate	Reoperation	Recurrence or residue
Zhao et al (2022) ¹¹ ,							
Total (N)	1231	1487	1372	1687	2,372	2,226	2,621
Pooled effect (95% CI); p value	MD -2.42 (-3.21 to -1.63);.0001	MD -0.23(-0.61 to 0.15);.60	MD -0.49 (-0.84 to -0.14);.006	MD -2.21 (-4.17 to -0.25);.03	OR 0.94 (0.67 to 1.32);.71	OR 1.67 (1.17 to 2.36);.004	OR 1.55 (1.07 to 2.24);.02
I² (p)	95%;.00001	51%;.03	90%;.00001	88%;.00001	0%;.65	0%;.89	0%;.93
Bai et al (2021) ⁹ ,							
Total (N)	NR	NR	NR	NR	NR		NR
Pooled effect (95% CI); p value	MD -2.59 (-3.87 to -1.31); <.001	MD 0.00 (-0.10 to 0.10);.991	MD -0.17 (-0.55 to 0.21);.384	MD -0.29 (-1.00 to 0.43);.434	relative risk 0.86 (0.63 to 1.18);.361		relative risk 1.65 (1.08 to 2.52);.021
I² (p)	72.1%;.001	0.0%;.996	88.3%; <.001	0.0%;.996	51.5%;.024		26.1%;.220
Gadjradj et al (2021) ¹⁰ ,							
Total (N)		621 and 152		621 and 152			
Pooled effect (95% CI)		3 to 6 month MD 0.05 (-0.10 to 0.21) 12 month MD 0.11 (-0.30 to 0.53)		3 to 6 month MD -0.09 (-0.24 to 0.07) 12 month MD -0.11 (-0.45 to 0.24)			
I² (p)		30%;.23		9%;.83			
Xu et al (2020) ⁸ ,							
Total (N)	NR	NR	NR	NR	NR	NR	NR
Pooled effect (95% CI); p value	OR -1.041 (-1.493 to -0.583);.000	6 months to 2 years OR -0.138 (-0.384 to 0.108);.270 2 years OR 0.020 (-0.193 to 0.233);.855	6 months to 2 years -0.456 (-0.947 to 0.034);.068 2 years OR -0.856 (-1.488 to -0.224);.008	6 months to 2 years -0.077 (-0.370 to 0.215);.604 2 years OR -0.425 (-0.724 to -0.127);.005	OR 0.972 (0.635 to 1.488);.896	OR 1.136 (0.415 to 3.108);.805	OR 1.306 (0.664 to 2.566);.439
I² (p)		53.8%;.090; 6 months to 2 years 4.4%;.351; 2 years	88%;.000; 6 months to 2 years 86.7%;.001; 2 years	75.3%;.000; 6 months to 2 years 52.7%;.121; 2 years			

Zhou et al (2020)⁷							
Total (N)						787	972
Pooled effect (95% CI); p value						OR 1.77 (1.18 to 2.64);.006	OR 1.60 (1.01 to 2.53);.05
I² (p)						0%;.97	0%;.94
Yu et al (2019)⁶							
Total (N)	707	NR	NR	NR	659		443
Pooled effect (95% CI); p value	MD -1.92 (-2.90 to -0.94); <.001	1 year postop or last follow-up: MD -0.07 (-0.22 to 0.08);.38	1 year postop or last follow-up: MD -0.41 (-0.76 to -0.06);.02	1 year postop or last follow-up: MD -0.27 (-1.71 to 1.16);.71	MD 1.01 (0.60 to 1.69);.98		MD 1.31 (0.54 to 3.17);.54
I² (p)	88%				0%		0%
Shi et al (2019)⁵							
Total (N)	1717	742	742	1337	1527	805	928
Pooled effect (95% CI); p value	MD -2.29 (3.03 to -1.55); <.00001	At last follow-up: MD -0.18 (-0.45 to 0.09);.19	At last follow-up: MD -0.77 (-1.31 to -0.24);.005	At last follow-up: MD -0.30 (-1.02 to 0.42);.41	OR 0.96 (0.65 to 1.43);.85	OR 2.67 (1.07 to 6.67);.04	OR 2.22 (1.02 to 4.83);.05
I² (p)	96%; <.00001	88%; <.00001	95%; <.00001	55%;.01	0%;.90	0%;.79	0%;.86
Phan et al (2017)⁴							
Total (N)	685	390		303	27,699	995	1081
Pooled effect (95% CI); p value	MD -4.79 (-6.52 to -3.07); <.00001	MD -0.04 (-0.37 to 0.30);.84		MD -1.88 (-4.06 to 0.29);.09	OR 0.77 (0.45 to 1.31);.33	OR 1.46 (0.33 to 6.43);.61	OR 1.12 (0.60 to 2.09);.73
I² (p)	99%; <.00001	70%;.003		67%;.03	60%;.004	66%;.004	0%;.97

CI: confidence interval; MD: mean difference; NR: not reported; ODI: Oswestry Disability Index; OR: odds ratio; RR: risk ratio; VAS: visual analogue scale; WMD: weighted mean difference

Results from the systematic reviews were fairly consistent with a significantly reduced length of hospitalization observed with endoscopic discectomy and sometimes significant improvements in VAS or ODI, but only at specific time points. Overall, no consistently significant improvement in VAS, ODI, total complication rate, reoperation, or recurrence was observed with endoscopic discectomy versus other interventions. Authors of the systematic reviews noted multiple limitations including the innate flaws of included studies (i.e., observational designs, a limited number of studies meeting criteria for inclusion, small sample sizes, lack of allocation concealment and blinding), different methodologies contributing to heterogeneity in analyses, loss of usable and sufficient data resulting in difficulty performing accurate analysis of

outcomes, and that a majority of the more recently completed studies were completed in China, which may affect the generalizability of the results to other populations.

Randomized Controlled Trials

More recent RCTs not included in any of the systematic reviews were also identified.^{87,88,89} Results of these trials are similar to those seen in the more comprehensive systematic reviews – percutaneous endoscopic discectomy was associated with a significant reduction in length of stay with no consistent or clinically meaningful improvements in patient-reported outcome measures such as VAS and ODI. Two of the 3 RCTs evaluated treatment-related morbidities, and reported a reduced incidence of intraoperative and postoperative complications and repeat surgeries with percutaneous endoscopic discectomy. Key characteristics, results, and limitations of these RCTs are summarized in the following tables.

Table 8. Characteristics of RCTs of Percutaneous Endoscopic Discectomy

Study	Countries	Sites	Dates	Participants	Interventions
Gadjraj et al 2022 ⁸⁷ ,	Netherlands	4	February 2016 to April 2019	Patients with sciatica caused by lumbar disc herniation	PTED vs microendoscopic discectomy
Ran et al 2021 ⁸⁸ ,	China	1	August 2016 to February 2020	Patients with highly migrated lumbar disc herniation	PELD with computerized tomography navigation vs open discectomy
Wang et al 2019 ⁸⁹ ,	China	1	July 2015 to July 2016	Patients with single-segment lumbar disc herniation with imaging results consistent with symptoms	PTED vs microendoscopic discectomy

PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials.

Table 9. Results of RCTs of Percutaneous Endoscopic Discectomy

Study	Length of stay (days)	Leg pain VAS	Lower back pain VAS	ODI	SF-36 PCS	Complication rates	Repeat surgery within 1 year
Gadjraj et al 2022 ⁸⁷ ,							
N	420	413	413	413	413	420	420

Pooled effect at 12 months (95% CI)	Median (IQR) PTED: 0 (0 to 0) Microendoscopic discectomy: 1 (1 to 1)	MD 7.1 (2.8 to 11.3)	MD 6 (2 to 10)	MD 5.3 (3.0 to 7.7)	MD -2.8 (-4.1 to -1.6)	PTED vs microendoscopic discectomy: Dural tears (n=0 vs 8) Nerve root injury (n=0 vs 1) Wound infection (n=3 vs 0) Cerebrospinal fluid leakage (n=1 vs 0)	PTED vs microendoscopic discectomy: n=9 (5%) vs 14 (6%)
p-value							
Ran et al 2021⁸⁸,							
N		66				66	
PELD with computerized tomography navigation at 12 months		0.58 ± 0.90				Infection, n=0 Recurrence, n=1	
Open discectomy at 12 months		0.75 ± 0.84				Infection, n=1 Recurrence, n=0	
p-value		.58				>.99	
N	90	90	90	90			
PTED	Postoperative: 3.01 ± 0.52	Preoperative mean score vs. 6 months after surgery: 7.21 vs. 1.05	Preoperative mean score vs 6 months after surgery: 6.40 vs. 1.36	Preoperative mean score vs 6 months after surgery: 58.21%			
				vs. 17.05%			
Microendoscopic discectomy	Postoperative: 6.68 ± 0.30	Preoperative mean score vs. 6 months after surgery: 7.09 vs. 0.98	Preoperative mean score vs 6 months after surgery: 6.34 vs. 1.65	Preoperative mean score vs 6 months after surgery: 57.17% vs. 16.98%			

P-value	.001	.097	.523	.864			
				2.6			

IQR: interquartile range; MD: mean difference; ODI: Oswestry Disability Index; PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials; SF-36 PCS: Short-Form-36 Physical Component Score; VAS: visual analogue scale.

Table 10. Study Relevance Limitations of the RCTs of Percutaneous Endoscopic Discectomy

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Gadraj et al 2022⁸⁷,	4. Limited to participants from 3 sites in the Netherlands				
Ran et al 2021⁸⁸,	4. Limited to participants from single site in China	4. PELD was used with computerized tomography navigation		1. Morbidity-related outcomes such as complications were limited	
Wang et al 2019⁸⁹,	4. Study population similar to other trials with regard to age, sex; however, included patients from a single Chinese hospital			1. Morbidity-related outcomes such as complication and reoperation rates were not reported	1,2. Outcomes reported only for 6 months of follow-up

PELD: percutaneous endoscopic lumbar discectomy; RCT: randomized controlled trials.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. ^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other. ^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest. ^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. ^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported. ^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 11. Study Design and Conduct Limitations of the RCTs of Percutaneous Endoscopic Discectomy

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Gadraj et al 2022⁸⁷,	4. A proportion of patients with a strong preference for PTED who were randomized to	1,2. Blinding did not occur				

	open microdiscectomy dropped out of the study after randomization					
Ran et al 2021⁸⁸,	3.Allocation concealment unclear	1,2. Blinding did not appear to occur			1. Power calculations not reported	
Wang et al 2019⁸⁹,	3.Allocation concealment unclear	1,2. Blinding did not appear to occur			1. Power calculations not reported	

PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials. The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. ^aAllocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias. ^bBlinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. ^cSelective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^dData Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). ^ePower key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference. ^fStatistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Observational Studies

Comparative observational studies with at least a 2-year follow-up are summarized below.

Yu et al (2021) published the results of a retrospective multicenter study that followed patients for 2 years after receipt of transforaminal percutaneous endoscopic discectomy (n=632) and microendoscopic discectomy (n=421) for lumbar disc herniation. Mean blood loss (p<.001) and mean duration of hospital stay (p=.018) were significantly reduced with transforaminal percutaneous endoscopic lumbar discectomy compared to microendoscopic discectomy. Rates of complications, recurrence, and revisions were similar in both groups. The VAS pain scores did not differ between groups after the first postoperative day. At 1 month postoperatively, there was a significant difference in ODI scores between groups (p=.016) in favor of transforaminal percutaneous endoscopic discectomy, but there was no significant difference at other time points.

Song et al (2021) published a retrospective single-center study that compared percutaneous endoscopic lumbar discectomy (n=306) and microendoscopic discectomy (n=116) in patients undergoing same day ambulatory surgery for lumbar disc herniation. Mean blood loss and mean duration of hospital stay were significantly less with percutaneous endoscopic lumbar discectomy (both p<.001 compared to microendoscopic discectomy). After 3 years of follow-up, e VAS pain scores for the back were also significantly lower in the percutaneous endoscopic lumbar discectomy group compared to the microendoscopic discectomy group (p=.001), but

there was no difference between groups in pain scores for the legs ($p=.224$). Overall recurrence rates ($p=.201$) and ODI scores ($p=.220$) were also similar between groups.

A number of observational studies have also assessed the learning curve^{90,91,92}, and the need for longer follow-up for endoscopic discectomy.^{93,94} The largest and longest follow-up to date has been reported by Choi et al (2015), who examined 10,228 patients at their institution who had had percutaneous endoscopic lumbar discectomy over a 12-year period.⁹⁵ They found that 4.3% of cases required reoperation in the first 6 weeks due to incomplete removal of herniated discs (2.8%), recurrence (0.8%), persistent pain (0.4%), and approach-related pain (0.2%).

Section Summary: Percutaneous Endoscopic Discectomy

The evidence for percutaneous endoscopic discectomy in individuals who have herniated intervertebral disc(s) includes a number of RCTs, systematic reviews, and comparative observational studies with at least 2 years of follow up. Many of the more recent RCTs are conducted at institutions within China. There are few reports from the United States. Overall, results from RCTs and systematic reviews reveal a significantly reduced length of hospitalization with endoscopic discectomy and occasionally significant improvements in VAS or ODI, but only at specific time points. No consistently significant improvement in VAS, ODI, total complication rate, reoperation, or recurrence was observed with percutaneous endoscopic discectomy versus other interventions.

Summary of Evidence

For individuals who have herniated intervertebral disc(s) who receive automated percutaneous discectomy, the evidence includes randomized controlled trials (RCTs) and systematic reviews of observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The published evidence from small RCTs is insufficient to evaluate the impact of automated percutaneous discectomy on the net health outcome. Well designed and executed RCTs are needed to determine the benefits and risks of this procedure.

The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have herniated intervertebral disc(s) who receive percutaneous endoscopic discectomy, the evidence includes a number of RCTs, systematic reviews, and observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. Many of the more recent RCTs are conducted at institutions within China. There are few reports from the United States. Results do not reveal a consistently significant improvement in patient-reported outcomes and treatment-related morbidity with percutaneous endoscopic discectomy in comparison to other discectomy interventions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2018 Input

Clinical input was sought to help determine whether the use of automated percutaneous discectomy or endoscopic percutaneous discectomy for individuals with herniated intervertebral discs would provide a clinically meaningful improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, clinical input was received from 3 respondents, including 2 specialty society-level response(s); no physician-level responses identified through a specialty society; 1 physician level response identified through an academic medical center. For individuals who have herniated intervertebral discs who receive automated percutaneous discectomy or percutaneous endoscopic discectomy, clinical input does not support a clinically meaningful improvement in net health outcome and does not indicate this use is consistent with generally accepted medical practice. Clinical input suggests that automated percutaneous discectomy may be an appropriate treatment option for the highly selected patient who has a small focal disc fragment compressing a lumbar nerve causing radiculopathy in the absence of lumbar stenosis or severe bony foraminal stenosis. Similarly, clinical input suggests that endoscopic percutaneous discectomy may be an appropriate treatment option for the highly selected patient who has a small focal disc herniation causing lumbar radiculopathy. However, respondents were mixed in the level of support for this indication, and overall the clinical input is not generally supportive of a clinically meaningful improvement in net health outcome.

2013 Input

In response to requests, input was received from 4 physician specialty societies and 3 academic medical centers while this policy was under review in 2013. Overall, input agreed that percutaneous and endoscopic discectomy are investigational. Most reviewers considered discectomy with tubular retractors to be a variant of open discectomy, with the only difference being the type of retraction used.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they are issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Institute for Health and Care Excellence

The NICE (2005) published guidance on automated percutaneous mechanical lumbar discectomy, indicating there was limited evidence of efficacy based on uncontrolled case series of heterogeneous groups of patients, and evidence from small RCTs showed conflicting results.⁹⁶ The guidance indicated that, in view of uncertainty about the efficacy of the procedure, it should not be done without special arrangements for consent and for audit or research. The guidance was considered for an update in 2009, but failed review criteria; the 2005 guidance is therefore considered current.

A NICE (2016) guidance on percutaneous transforaminal endoscopic lumbar discectomy for sciatica was published.⁹⁷ The guidance stated that current evidence is adequate to support the use of percutaneous transforaminal endoscopic lumbar discectomy for sciatica. Choice of operative procedure (open discectomy, microdiscectomy, or percutaneous endoscopic approaches) may be influenced by symptoms, location, and size of the prolapsed disc.

A NICE (2016) guidance on percutaneous interlaminar endoscopic lumbar discectomy for sciatica was also published.⁹⁸ The guidance stated that current evidence is adequate to support the use of percutaneous interlaminar endoscopic lumbar discectomy for sciatica. Choice of operative procedure (open discectomy, microdiscectomy, or percutaneous endoscopic approaches) may be influenced by symptoms, location, and size of the prolapsed disc.

American Society of Interventional Pain Physicians

The guidelines from the American Society of Interventional Pain Physicians (2013) indicated that the evidence for percutaneous disc decompression with the Dekompessor was limited.³ There were no recommended indications for the Dekompessor.

North American Spine Society

The North American Spine Society (2014) published clinical guidelines on the diagnosis and treatment of lumbar disc herniation⁹⁹, Table 12 summarizes recommendations specific to percutaneous endoscopic discectomy and automated percutaneous discectomy.

Table 12. Recommendations for Lumbar Disc Herniation With Radiculopathy

Recommendations	Grade or LOE ^a
Endoscopic percutaneous discectomy is suggested for carefully selected patients to reduce early postoperative disability and reduce opioid use compared with open discectomy.	B
There is insufficient evidence to make a recommendation for or against the use of automated percutaneous discectomy compared with open discectomy.	I
Endoscopic percutaneous discectomy may be considered for treatment.	C
Automated percutaneous discectomy may be considered for treatment.	C

Patients undergoing percutaneous endoscopic discectomy experience better outcomes if <40 years and symptom duration <3 months.

II

LOE: level of evidence.

^a Grade B: fair evidence (level II or III studies with consistent findings; grade C: poor quality evidence (level IV or V studies). Level of evidence II:

lesser quality randomized controlled trial (e.g., <80% follow-up, no blinding, or improper randomization), prospective comparative study, systematic review of level II studies or level I studies with inconsistent results; level of evidence III: case control, retrospective, systematic review of level III studies; level of evidence IV: case series; level of evidence V: expert opinion.

American Pain Society

The clinical practice guidelines from the American Pain Society (2009) found insufficient evidence to evaluate alternative surgical methods to standard open discectomy and microdiscectomy, including laser or endoscopic-assisted techniques, various percutaneous techniques, coblation nucleoplasty, or the Dekompressor.¹⁰⁰

American Society of Pain and Neuroscience

The American Society of Pain and Neuroscience (ASPN; 2022) published clinical guidance for interventional treatments for low back pain.[Sayed D, Grider J, Strand N, et al. The American S.... ; 15: 3729-3832. PMID 36510616] The guideline states that discectomy procedures (such as percutaneous and endoscopic disc procedures) have favorable safety and efficacy profiles for the treatment of lumbar disc herniation with persistent radicular symptoms; however, it is stated that further research is needed to evaluate complications rates in order for these procedures to supplant classic open microdiscectomy. Recommendations specific to percutaneous endoscopic discectomy are summarized in Table 13.

National Comprehensive Cancer Network

The National Comprehensive Cancer Network (NCCN) clinical practice guidelines for central nervous system cancers (v.1.2023) states that MRgLITT “may be considered for patients who are poor surgical candidates (craniotomy or resection). Potential indications include relapsed brain metastases, radiation necrosis, and recurrent glioblastoma.” (Category 2B)³⁸,The guidelines additionally state that LITT “can be considered on a case-by-case basis for treatment of radiation necrosis in patients with a history of RT [radiation therapy] for primary brain tumor or metastatic disease. Consultation with adept neurosurgeons trained in LITT should be done when the procedure is considered.”

Table 13. Recommendations for Percutaneous and Endoscopic Procedures

Recommendation	Grade ^a	Level of Evidence ^b	Level of Certainty [Net Benefit] ^c
Percutaneous Endoscopic Discectomy	B	I-a	High

^a Grade B: (The ASPN Back Group recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial..

^b Evidence Level: I-A: At least one controlled and randomized clinical trial, properly designed

U.S. Preventative Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Currently unpublished trials that might influence this review are listed in Table 14.

Table 14. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT01997086	Percutaneous Transforaminal Endoscopic Discectomy (PTED) vs. Microendoscopic Discectomy (MED) for the treatment of Lumbar Disc Herniation: A Prospective Randomized Controlled Study	125	Aug 2023
NCT02602093	Percutaneous Transforaminal Endoscopic Discectomy vs. Open Microdiscectomy for Lumbar Disc Herniation (PTED-study)	682	May 2024
Unpublished			
NCT02742311	EuroPainClinics® Study V Prospective Observational Study (EPCSV)	500	Dec 2021

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Essential Health Benefits

The Affordable Care Act (ACA) requires fully insured non-grandfathered individual and small group benefit plans to provide coverage for ten categories of Essential Health Benefits (“EHBs”), whether the benefit plans are offered through an Exchange or not. States can define EHBs for their respective state.

States vary on how they define the term small group. In Idaho, a small group employer is defined as an employer with at least two but no more than fifty eligible employees on the first day of the plan or contract year, the majority of whom are employed in Idaho. Large group employers, whether they are self-funded or fully insured, are not required to offer EHBs, but may voluntarily offer them.

The ACA requires any benefit plan offering EHBs to remove all dollar limits for EHBs.

Applicable Coding

CPT Category III Codes – Not Covered

CPT Codes	Number	Description
-----------	--------	-------------

	62287	Decompression procedure, percutaneous, of nucleus pulposus of intervertebral disc, any method, utilizing needle-based technique to remove disc material under fluoroscopic imaging or other form of indirect visualization, with discography and/or epidural injection(s) at the treated level(s), when performed, single or multiple levels, lumbar
	62380	Endoscopic decompression of spinal cord, nerve root(s), including laminotomy, partial facetectomy, foraminotomy, discectomy and/or excision of herniated intervertebral disc, 1 interspace, lumbar
	0274T	Percutaneous laminotomy/laminectomy (interlaminar approach) for decompression of neural elements, (with or without ligamentous resection, discectomy, facetectomy and/or foraminotomy), any method, under indirect
		image guidance (e.g., fluoroscopic, CT), single or multiple levels, unilateral or bilateral;cervical or thoracic
	0275T	;lumbar
HCPCS	C2614	Probe, percutaneous lumbar discectomy
ICD-10-CM	C71.0c71.9	Investigational for all relevant diagnoses
ICD-10-PCS		ICD-10-PCS codes are only used for inpatient services
	0S523ZZ 0S524ZZ	Lower joints, destruction, lumbar vertebral disc, no device, codes for percutaneous or percutaneous endoscopic
	0S543ZZ 0S544ZZ	Destruction lumbosacral disc, no device, codes for percutaneous or percutaneous endoscopic
	0R533ZZ 0R534ZZ	Upper joints, destruction, cervical vertebral disc, no device, codes for percutaneous or percutaneous endoscopic
	0R593ZZ 0R594ZZ	Upper joints, destruction, thoracic vertebral disc, no device, codes for percutaneous or percutaneous endoscopic
	0R5B3ZZ 0R5B4ZZ	Upper joints, destruction, thoracolumbar vertebral disc, no device, codes for percutaneous or percutaneous endoscopic

Type of Service	Surgery	

Vendors
<ul style="list-style-type: none"> • MedCom • Health Plan Services

References
<ol style="list-style-type: none"> 1. Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. <i>Spine J.</i> Jun 01 2015; 15(6): 1461-77. PMID 24412033 2. Haines SJ, Jordan N, Boen JR, et al. Discectomy strategies for lumbar disc herniation: results of the LAPDOG trial. <i>J Clin Neurosci.</i> Jul 2002; 9(4): 411-7. PMID 12217670 3. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. <i>Pain Physician.</i> Apr 2013; 16(2 Suppl): S49-283. PMID 23615883 4. Phan K, Xu J, Schultz K, et al. Full-endoscopic versus micro-endoscopic and open discectomy: A systematic review and meta-analysis of outcomes and complications. <i>Clin Neurol Neurosurg.</i> Mar 2017; 154: 1-12. PMID 28086154 5. Shi R, Wang F, Hong X, et al. Comparison of percutaneous endoscopic lumbar discectomy versus microendoscopic discectomy for the treatment of lumbar disc herniation: a meta-analysis. <i>Int Orthop.</i> Apr 2019; 43(4): 923-937. PMID 30547214 6. Yu P, Qiang H, Zhou J, et al. Percutaneous Transforaminal Endoscopic Discectomy versus Micro-Endoscopic Discectomy for Lumbar Disc Herniation. <i>Med Sci Monit.</i> Mar 30 2019; 25: 2320-2328. PMID 30927349 7. Zhao XM, Yuan QL, Liu L, et al. Is It Possible to Replace Microendoscopic Discectomy with Percutaneous Transforaminal Discectomy for Treatment of Lumbar Disc Herniation? A Meta-Analysis Based on Recurrence and Revision Rate. <i>J Korean Neurosurg Soc.</i> Jul 2020; 63(4): 477-486. PMID 32380585 8. Xu J, Li Y, Wang B, et al. Minimum 2-Year Efficacy of Percutaneous Endoscopic Lumbar Discectomy versus Microendoscopic Discectomy: A Meta-Analysis. <i>World Neurosurg.</i> Jun 2020; 138: 19-26. PMID 32109644 9. Bai X, Lian Y, Wang J, et al. Percutaneous endoscopic lumbar discectomy compared with other surgeries for lumbar disc herniation: A meta-analysis. <i>Medicine (Baltimore).</i> Mar 05 2021; 100(9): e24747. PMID 33655938 10. Gadjradj PS, Harhangi BS, Amelink J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus Open Microdiscectomy for Lumbar Disc Herniation: A Systematic Review and Meta-analysis. <i>Spine (Phila Pa 1976).</i> Apr 15 2021; 46(8): 538-549. PMID 33290374 11. Zhao XM, Chen AF, Lou XX, et al. Comparison of Three Common Intervertebral Disc Discectomies in the Treatment of Lumbar Disc Herniation: A Systematic Review and Meta-Analysis Based on Multiple Data. <i>J Clin Med.</i> Nov 08 2022; 11(22). PMID 36431083 12. Ma C, Li H, Zhang T, et al. Comparison of Percutaneous Endoscopic Interlaminar Discectomy and Open Fenestration Discectomy for Single-Segment Huge Lumbar Disc Herniation: A Two-year Follow-up Retrospective Study. <i>J Pain Res.</i> 2022; 15: 1061-1070. PMID 35444463

13. Wang SF, Hung SF, Tsai TT, et al. Better Functional Outcome and Pain Relief in the Far-Lateral-Outside-in Percutaneous Endoscopic Transforaminal Discectomy. *J Pain Res.* 2021; 14: 3927-3934. PMID 35002312
14. Rajamani PA, Goparaju P, Kulkarni AG, et al. A 2-Year Outcomes and Complications of Various Techniques of Lumbar Discectomy: A Multicentric Retrospective Study. *World Neurosurg.* Dec 2021; 156: e319-e328. PMID 34555576
15. Jing Z, Li L, Song J. Percutaneous transforaminal endoscopic discectomy versus microendoscopic discectomy for upper lumbar disc herniation: a retrospective comparative study. *Am J Transl Res.* 2021; 13(4): 3111-3119. PMID 34017479
16. Jarebi M, Awaf A, Lefranc M, et al. A matched comparison of outcomes between percutaneous endoscopic lumbar discectomy and open lumbar microdiscectomy for the treatment of lumbar disc herniation: a 2-year retrospective cohort study. *Spine J.* Jan 2021; 21(1): 114-121. PMID 32683107
17. Meyer G, DA Rocha ID, Cristante AF, et al. Percutaneous Endoscopic Lumbar Discectomy Versus Microdiscectomy for the Treatment of Lumbar Disc Herniation: Pain, Disability, and Complication Rate-A Randomized Clinical Trial. *Int J Spine Surg.* Feb 2020; 14(1): 72-78. PMID 32128306
18. Chen Z, Zhang L, Dong J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus Microendoscopic Discectomy for Lumbar Disc Herniation: Two-Year Results of a Randomized Controlled Trial. *Spine (Phila Pa 1976).* Apr 15 2020; 45(8): 493-503. PMID 31703056
19. Kim CH, Chung CK, Choi Y, et al. The Long-term Reoperation Rate Following Surgery for Lumbar Herniated Intervertebral Disc Disease: A Nationwide Sample Cohort Study With a 10-year Follow-up. *Spine (Phila Pa 1976).* Oct 01 2019; 44(19): 1382-1389. PMID 30973508
20. Ahn Y, Lee SG, Son S, et al. Transforaminal Endoscopic Lumbar Discectomy Versus Open Lumbar Microdiscectomy: A Comparative Cohort Study with a 5-Year Follow-Up. *Pain Physician.* May 2019; 22(3): 295-304. PMID 31151337
21. Liu X, Yuan S, Tian Y, et al. Comparison of percutaneous endoscopic transforaminal discectomy, microendoscopic discectomy, and microdiscectomy for symptomatic lumbar disc herniation: minimum 2-year follow-up results. *J Neurosurg Spine.* Mar 2018; 28(3): 317-325. PMID 29303471
22. Sun Y, Zhang W, Qie S, et al. Comprehensive comparing percutaneous endoscopic lumbar discectomy with posterior lumbar internal fixation for treatment of adjacent segment lumbar disc prolapse with stable retrolisthesis: A retrospective case-control study. *Medicine (Baltimore).* Jul 2017; 96(29): e7471. PMID 28723757
23. Jeong, J.S.; Lee, S.H.; Lee, S.J.; Hwang, B.W. The Clinical Comparison between Open Surgery and Percutaneous Endoscopic Lumbar Discectomy in Extraforaminal Lumbar Disc Herniation. *J. Korean Neurosurg. Soc.* 2006, 39, 413418.
24. Akçakaya MO, Yörükoğlu AG, Aydoseli A, et al. Serum creatine phosphokinase levels as an indicator of muscle injury following lumbar disc surgery: Comparison of fully endoscopic discectomy and microdiscectomy. *Clin Neurol Neurosurg.* Jun 2016; 145: 74-8. PMID 27101087
25. Choi KC, Shim HK, Hwang JS, et al. Comparison of Surgical Invasiveness Between Microdiscectomy and 3 Different Endoscopic Discectomy Techniques for Lumbar Disc Herniation. *World Neurosurg.* Aug 2018; 116: e750-e758. PMID 29787880

26. Dai HJ, Zhang X, Wang LT, et al. The effect of percutaneous transforaminal endoscopic discectomy (PTED) on serum inflammatory factors and pain in patients with lumbar disc herniation after surgery. *Int J Clin Exp Med* 2020;13:597603.
27. Krappel FA, Schmitz R, Bauer E, et al. Open or endoscopic nucleotomy?. *Orthopadische Praxis* 2001;37:1649.
28. Tacconi L, Giordan E. Endoscopic transforaminal discectomy vs. far lateral discectomy for extraforaminal disc protrusions: our experience. *NeuroQuantology* 2019;17:1822.
29. Tacconi L, Signorelli F, Giordan E. Is Full Endoscopic Lumbar Discectomy Less Invasive Than Conventional Surgery? A Randomized MRI Study. *World Neurosurg.* Jun 2020; 138: e867-e875. PMID 32251813
30. Tao XZ, Jing L, Li JH. Therapeutic effect of transforaminal endoscopic spine system in the treatment of prolapse of lumbar intervertebral disc. *Eur Rev Med Pharmacol Sci.* Jul 2018; 22(1 Suppl): 103-110. PMID 30004561
31. Wang H, Song Y, Cai L. Effect of percutaneous transforaminal lumbar spine endoscopic discectomy on lumbar disc herniation and its influence on indexes of oxidative stress. *Biomed Res* 2017;28:. Applicable Lines of Business: Individual Qualified Health Plan Medicare Supplement Large Group Small Group Original Policy Date _07/01/2024) Review Cycle: (Annually) Document Retention: 10 years Proprietary and Confidential 36 | Page
32. Xu G, Zhang C, Zhu K, et al. Endoscopic removal of nucleus pulposus of intervertebral disc on lumbar intervertebral disc protrusion and the influence on inflammatory factors and immune function. *Exp Ther Med.* Jan 2020; 19(1): 301-307. PMID 31853303
33. Ahn SS, Kim SH, Kim DW, et al. Comparison of Outcomes of Percutaneous Endoscopic Lumbar Discectomy and Open Lumbar Microdiscectomy for Young Adults: A Retrospective Matched Cohort Study. *World Neurosurg.* Feb 2016; 86: 250-8. PMID 26409086
34. Chang F, Zhang T, Gao G, et al. Therapeutic effect of percutaneous endoscopic lumbar discectomy on lumbar disc herniation and its effect on oxidative stress in patients with lumbar disc herniation. *Exp Ther Med.* Jan 2018; 15(1): 295-299. PMID 29250152
35. Liu C, Zhou Y. Percutaneous Endoscopic Lumbar Discectomy and Minimally Invasive Transforaminal Lumbar Interbody Fusion for Recurrent Lumbar Disk Herniation. *World Neurosurg.* Feb 2017; 98: 14-20. PMID 27773858
36. Pan Z, Ha Y, Yi S, et al. Efficacy of Transforaminal Endoscopic Spine System (TESSYS) Technique in Treating Lumbar Disc Herniation. *Med Sci Monit.* Feb 18 2016; 22: 530-9. PMID 26887645
37. Yao Y, Zhang H, Wu J, et al. Comparison of Three Minimally Invasive Spine Surgery Methods for Revision Surgery for Recurrent Herniation After Percutaneous Endoscopic Lumbar Discectomy. *World Neurosurg.* Apr 2017; 100: 641-647.e1. PMID 28153616
38. Yao Y, Zhang H, Wu J, et al. Minimally Invasive Transforaminal Lumbar Interbody Fusion Versus Percutaneous Endoscopic Lumbar Discectomy: Revision Surgery for Recurrent Herniation After Microendoscopic Discectomy. *World Neurosurg.* Mar 2017; 99: 89-95. PMID 27919762
39. Gibson JNA, Subramanian AS, Scott CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. *Eur Spine J.* Mar 2017; 26(3): 847-856. PMID 27885470
40. Hsu HT, Chang SJ, Yang SS, et al. Learning curve of full-endoscopic lumbar discectomy. *Eur Spine J.* Apr 2013; 22(4): 727-33. PMID 23076645

41. Kim MJ, Lee SH, Jung ES, et al. Targeted percutaneous transforaminal endoscopic discectomy in 295 patients: comparison with results of microscopic discectomy. *Surg Neurol.* Dec 2007; 68(6): 623-631. PMID 18053857
42. Qu JX, Li QZ, Chen M : Comparative study of PTED and MED for monosegmentnlumbar disc herniation. *Chin J Bone Joint Inj* 32 : 70-71,2017
43. Wang H, Cheng J, Xiao H, et al. Adolescent lumbar disc herniation: experience from a large minimally invasive treatment centre for lumbar degenerative disease in Chongqing, China. *Clin Neurol Neurosurg.* Aug 2013; 115(8): 1415-9. PMID 23419406
44. Zhao W, Li CQ, Zhou Y, Wang J, Zheng WJ : Surgical treatment of thelumbar disc herniated discs using transforaminal endoscopic surgeryssystem. *Orthop J China* 20 : 1191-1195, 2012
45. Yoon SM, Ahn SS, Kim KH, et al. Comparative Study of the Outcomes of Percutaneous Endoscopic Lumbar Discectomy and Microscopic Lumbar Discectomy Using the Tubular Retractor System Based on the VAS, ODI, and SF-36. *Korean J Spine.* Sep 2012; 9(3): 215-22. PMID 25983818
46. Li M, Yang H, Yang Q. Full-Endoscopic Technique Discectomy Versus Microendoscopic Discectomy for the Surgical Treatment of Lumbar Disc Herniation. *Pain Physician.* 2015; 18(4): 359-63. PMID 26218939
47. Sinkemani A, Hong X, Gao ZX, et al. Outcomes of Microendoscopic Discectomy and Percutaneous Transforaminal Endoscopic Discectomy for the Treatment of Lumbar Disc Herniation: A Comparative Retrospective Study. *Asian Spine J.* Dec 2015; 9(6): 833-40. PMID 26713113
- Applicable Lines of Business: Individual Qualified Health Plan Medicare Supplement Large Group Small Group Original Policy Date _07/01/2024) Review Cycle: (Annually) Document Retention: 10 years Proprietary and Confidential 37 | P a g e
48. Song HP, Sheng HF, Xu WX. A case-control study on the treatment of protrusion of lumbar intervertebral disc through PELD and MED. *Exp Ther Med.* Oct 2017; 14(4): 3708-3712. PMID 29042967
49. Tu Z, Li YW, Wang B, et al. Clinical Outcome of Full-endoscopic Interlaminar Discectomy for Single-level Lumbar Disc Herniation: A Minimum of 5-year Follow-up. *Pain Physician.* Mar 2017; 20(3): E425-E430. PMID 28339442
50. Li H, Jiang C, Mu X, et al. Comparison of MED and PELD in the Treatment of Adolescent Lumbar Disc Herniation: A 5-Year Retrospective Follow-Up. *World Neurosurg.* Apr 2018; 112: e255-e260. PMID 29325949
51. Abudurexiti T, Qi L, Muheremu A, et al. Micro-endoscopic discectomy versus percutaneous endoscopic surgery for lumbar disk herniation. *J Int Med Res.* Sep 2018; 46(9): 3910-3917. PMID 29900752
52. Chen Z, Zhang L, Dong J, et al. Percutaneous transforaminal endoscopic discectomy compared with microendoscopic discectomy for lumbar disc herniation: 1-year results of an ongoing randomized controlled trial. *J Neurosurg Spine.* Mar 2018; 28(3): 300-310. PMID 29303469
53. Liu T, Zhou Y, Wang J, et al. Clinical efficacy of three different minimally invasive procedures for far lateral lumbar disc herniation. *Chin Med J (Engl).* Mar 2012; 125(6): 1082-8. PMID 22613535
54. Wu XC, Zhou Y, Li CQ. Percutaneous tranforaminal endoscopic discectomy versus microendoscopic discectomy for lumbar disc herniation: a prospective randomized controlled study. *J Third Mil Med Univ.* 2009;31(9):843-846.

55. Yang L, Liao XQ, Zhao XJ, et al. Comparison of surgical outcomes between percutaneous transforaminal endoscopic discectomy and micro-endoscopic discectomy for lumbar disc herniation. *China J Endosc.* 2015;21(9):962-965
56. Duan XF, Jin W, Chen JJ, et al. Contrast observation of comparing microendoscopic discectomy with percutaneous endoscopic lumbar discectomy for the treatment of simple lumbar disc herniation. *Chin J Clin.* 2016;10(1):144-147
57. Zhao XW, Han K, Ji ZW, et al. Comparison of efficacy between microendoscopic discectomy and percutaneous endoscopic lumbar discectomy for treatment of lumbar disc herniation. *Prog Mod Biomed.* 2016;16(23):4454-4457
58. Ding YZ, Hu JN, Zhou Y, et al. Study on the effect contrast between microendoscopic discectomy and percutaneous endoscopic lumbar discectomy using TESSYS technique for the treatment of lumbar disc herniation. *J Cervicodynia & Lumbodynia.* 2017;38(5):492-493
59. Li ZY, Guo PG, Han D, et al. Analysis of curative effects and prognosis in different procedures of discectomy for patients with lumbar disc herniation. *J Clin Med Pract.* 2017;21(15):149-150,158
60. Liu HP, Hao DJ, Wang XD, et al. Comparison of two surgeries in treatment of lumbar disc herniation. *Chin J Pain Med.* 2017;23(6):438-442
61. Luo DK, Zhou NX, Zhao HW, et al. Clinical effectiveness of minimally invasive treatment for lumbar disc herniation. *Orthopaedics.* 2017;8(6):439-444
62. Qu JX, Li QZ, Chem M, et al. Comparison of the efficacies between percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy for the treatment of single-segmental lumbar disc herniation. *Chin J Bone Jt Inj.* 2017;32(1):70-71
63. Chen Q, Qin L, Li MW, et al. Comparison of the therapeutic effect of percutaneous transforaminal endoscopic discectomy and posterior discectomy on senile single segmental lumbar disc herniation. *Chin J Front Med Sci.* 2018;10(2):60-64
64. Wu YM, Bai M, Yin HP, et al. Comparison of the efficacies between two kinds of minimally invasive procedures for the treatment of simple lumbar disc herniation. *J Pract Orthop.* 2018;24(4):357-360
65. Belykh E, Giers MB, Preul MC, et al. Prospective Comparison of Microsurgical, Tubular-Based Endoscopic, and Endoscopically Assisted Discectomies: Clinical Effectiveness and Complications in Railway Workers. *World Neurosurg.* Jun 2016; 90: 273-280. PMID 26898494
66. Chen HC, Lee CH, Wei L, et al. Comparison of percutaneous endoscopic lumbar discectomy and open lumbar surgery for adjacent segment degeneration and recurrent disc herniation. *Neurol Res Int.* 2015; 2015: 791943. PMID 25861474
67. Choi KC, Kim JS, Park CK. Percutaneous Endoscopic Lumbar Discectomy as an Alternative to Open Lumbar Microdiscectomy for Large Lumbar Disc Herniation. *Pain Physician.* Feb 2016; 19(2): E291-300. PMID 26815256
68. Garg B, Nagraja UB, Jayaswal A. Microendoscopic versus open discectomy for lumbar disc herniation: a prospective randomised study. *J Orthop Surg (Hong Kong).* Apr 2011; 19(1): 30-4. PMID 21519072
69. Hermantin FU, Peters T, Quartararo L, et al. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am.* Jul 1999; 81(7): 958-65. PMID 10428127

70. Huang TJ, Hsu RW, Li YY, et al. Less systemic cytokine response in patients following microendoscopic versus open lumbar discectomy. *J Orthop Res.* Mar 2005; 23(2): 406-11. PMID 15734255
 71. Hussein M, Abdeldayem A, Mattar MM. Surgical technique and effectiveness of microendoscopic discectomy for large uncontained lumbar disc herniations: a prospective, randomized, controlled study with 8 years of follow-up. *Eur Spine J.* Sep 2014; 23(9): 1992-9. PMID 24736930
 72. Kleinpeter G, Markowitsch MM, Böck F. Percutaneous endoscopic lumbar discectomy: minimally invasive, but perhaps only minimally useful?. *Surg Neurol.* Jun 1995; 43(6): 534-9; discussion 540-1. PMID 7482230
 73. Lee DY, Shim CS, Ahn Y, et al. Comparison of percutaneous endoscopic lumbar discectomy and open lumbar microdiscectomy for recurrent disc herniation. *J Korean Neurosurg Soc.* Dec 2009; 46(6): 515-21. PMID 20062565
 74. Martín-Láez R, Martínez-Agüeros JA, Suárez-Fernández D, et al. Complications of endoscopic microdiscectomy using the EASYGO! system: is there any difference with conventional discectomy during the learning-curve period?. *Acta Neurochir (Wien).* Jun 2012; 154(6): 1023-32. PMID 22446750
 75. Mayer HM, Brock M. Percutaneous endoscopic discectomy: surgical technique and preliminary results compared to microsurgical discectomy. *J Neurosurg.* Feb 1993; 78(2): 216-25. PMID 8267686
 76. Ohya J, Oshima Y, Chikuda H, et al. Does the microendoscopic technique reduce mortality and major complications in patients undergoing lumbar discectomy? A propensity score-matched analysis using a nationwide administrative database. *Neurosurg Focus.* Feb 2016; 40(2): E5. PMID 26828886
 77. Pan L, Zhang P, Yin Q. Comparison of tissue damages caused by endoscopic lumbar discectomy and traditional lumbar discectomy: a randomised controlled trial. *Int J Surg.* 2014; 12(5): 534-7. PMID 24583364
 78. Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. *Neurosurgery.* Sep 2007; 61(3): 545-9; discussion 549. PMID 17881967
 79. Ruetten S, Komp M, Merk H, et al. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine (Phila Pa 1976).* Apr 20 2008; 33(9): 931-9. PMID 18427312
 80. Ruetten S, Komp M, Merk H, et al. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision. *J Spinal Disord Tech.* Apr 2009; 22(2): 122-9. PMID 19342934
 81. Sasaoka R, Nakamura H, Konishi S, et al. Objective assessment of reduced invasiveness in MED. Compared with conventional one-level laminotomy. *Eur Spine J.* May 2006; 15(5): 577-82. PMID 15926058
 82. Schizas C, Tsiridis E, Saksena J. Microendoscopic discectomy compared with standard microsurgical discectomy for treatment of uncontained or large contained disc herniations. *Neurosurgery.* Oct 2005; 57(4 Suppl): 357-60; discussion 357-60. PMID 16234685
- Applicable Lines of Business: Individual Qualified Health Plan Medicare Supplement Large Group Small Group Original Policy Date _07/01/2024) Review Cycle: (Annually) Document Retention: 10 years Proprietary and Confidential 39 | P a g e

83. Teli M, Lovi A, Brayda-Bruno M, et al. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy. *Eur Spine J*. Mar 2010; 19(3): 443-50. PMID 20127495
84. Ruetten S, Komp M, Merk H, et al. Use of newly developed instruments and endoscopes: full-endoscopic resection of lumbar disc herniations via the interlaminar and lateral transforaminal approach. *J Neurosurg Spine*. Jun 2007; 6(6): 521-30. PMID 17561740
85. Ruetten S, Komp M, Merk H, et al. Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9-mm endoscopes: a prospective, randomized, controlled study. *Spine (Phila Pa 1976)*. Apr 20 2008; 33(9): 940-8. PMID 18427313
86. Lee SH, Chung SE, Ahn Y, et al. Comparative radiologic evaluation of percutaneous endoscopic lumbar discectomy and open microdiscectomy: a matched cohort analysis. *Mt Sinai J Med*. Sep 2006; 73(5): 795-801. PMID 17008941
87. Gadraj PS, Rubinstein SM, Peul WC, et al. Full endoscopic versus open discectomy for sciatica: randomised controlled non-inferiority trial. *BMJ*. Feb 21 2022; 376: e065846. PMID 35190388
88. Ran B, Wei J, Yang J, et al. Quantitative Evaluation of the Trauma of CT Navigation PELD and OD in the Treatment of HLDH: A Randomized, Controlled Study. *Pain Physician*. Jul 2021; 24(4): E433-E441. PMID 34213868
89. Wang F, Guo D, Sun T, et al. A comparative study on short-term therapeutic effects of percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy on lumbar disc herniation. *Pak J Med Sci*. 2019; 35(2): 426-431. PMID 31086527
90. Lee DY, Lee SH. Learning curve for percutaneous endoscopic lumbar discectomy. *Neurol Med Chir (Tokyo)*. Sep 2008; 48(9): 383-8; discussion 388-9. PMID 18812679
91. Wang B, Lü G, Patel AA, et al. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. *Spine J*. Feb 2011; 11(2): 122-30. PMID 21296295
92. Tenenbaum S, Arzi H, Herman A, et al. Percutaneous Posterolateral Transforaminal Endoscopic Discectomy: Clinical Outcome, Complications, and Learning Curve Evaluation. *Surg Technol Int*. Dec 2011; 21: 278-83. PMID 22505002
93. Casal-Moro R, Castro-Menéndez M, Hernández-Blanco M, et al. Long-term outcome after microendoscopic discectomy for lumbar disk herniation: a prospective clinical study with a 5-year follow-up. *Neurosurgery*. Jun 2011; 68(6): 1568-75; discussion 1575. PMID 21311384
94. Wang M, Zhou Y, Wang J, et al. A 10-year follow-up study on long-term clinical outcomes of lumbar microendoscopic discectomy. *J Neurol Surg A Cent Eur Neurosurg*. Aug 2012; 73(4): 195-8. PMID 22825836
95. Choi KC, Lee JH, Kim JS, et al. Unsuccessful percutaneous endoscopic lumbar discectomy: a single-center experience of 10,228 cases. *Neurosurgery*. Apr 2015; 76(4): 372-80; discussion 380-1; quiz 381. PMID 25599214
96. National Institute for Health and Care Excellence (NICE). Automated percutaneous mechanical lumbar discectomy guidance [IPG141]. 2005; <http://guidance.nice.org.uk/IPG141/Guidance/pdf/English>. Accessed May 8, 2023.
97. National Institute for Health and Care Excellence (NICE). Percutaneous interlaminar endoscopic lumbar discectomy for sciatica [IPG555]. 2016; <https://www.nice.org.uk/guidance/ipg555>. Accessed May 7, 2023.
National Institute for Health and Care Excellence (NICE). Percutaneous transforaminal endoscopic lumbar discectomy for sciatica [IPG556]. 2016; <https://www.nice.org.uk/guidance/ipg556>. Accessed May 6, 2023.

98. Kreiner DS, Hwang SW, Easa JE, et al. An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. *Spine J.* Jan 2014; 14(1): 180-91. PMID 24239490
99. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976).* May 01 2009; 34(10): 1066-77. PMID 19363457
 Applicable Lines of Business: Individual Qualified Health Plan Medicare Supplement Large Group Small Group Original Policy Date _07/01/2024) Review Cycle: (Annually) Document Retention: 10 years Proprietary and Confidential 40 | Page
 Applicable Lines of Business: Individual Qualified Health Plan Original Policy Date _07/01/2024) Medicare Supplement Review Cycle: (Annually) Large Group Document Retention: 10 years Small Group Proprietary and Confidential 41 | Page
100. Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPEN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain. *J Pain Res.* 2022; 15: 3729-3832. PMID 36510616

Review/Revision/Approval History	
Date	Description
07/01/2024	New Policy
04/27/2026	Reviewed and Approved by Policy Committee

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult appropriate healthcare providers for medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member’s benefit plan, effective when services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Including or excluding a procedure, diagnosis, or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as they apply to an individual member.

Mountain Health Co-Op makes no representations and accepts no liability regarding the content of any external information cited or relied upon in this policy. Mountain Health Co-Op updates its Coverage Policies regularly and reserve

es the right to amend these policies and give notice per State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Mountain Health Co-Op.

”Mountain Health Co-Op” and its accompanying logo and marks are protected and registered trademarks of Mountain Health Co-Op. The content of this Service is proprietary and protected by

copyright. You may access the copyrighted content of this Service only for purposes outlined in these Conditions of Use.

© CPT Only – American Medical Association