

Policy	MM-089
Effective Date	09/01/2024
Reviewed/Revised Date	04/27/2026
Next Review Date	04/27/2027
Origination Date	07/01/2024
Originated Department	Clinical Operations

Computer Assisted Navigation for Orthopedic Procedure

Audience
Medical Management

Purpose
<p>Medical policies provide general support for applying Mountain Health Co-Op member policy document coverage decisions and must reference the member-specific benefit plan document. The terms of the member-specific Policy document may differ from the standard benefit plan on which this medical policy is based. If there is a conflict between a member-specific policy document and the Mountain Health Co-Op medical policy, the member-specific policy document supersedes this medical policy. Any person(s) applying this medical policy must identify member eligibility, the member-specific policy document, and related policies or guidelines before applying this medical policy, including the existence of any state or federal guidance. Mountain Health Co-Op medical policies are designed for informational purposes only and are not an authorization, explanation of benefits, or contract. Receipt of benefits is subject to satisfaction of all terms and conditions of the member-specific policy document coverage. Mountain Health Co-Op reserves the sole discretionary right to modify all policies and guidelines at any time.</p>

Definition
N/A

Policy/Procedure
<p>Policy Statement and Criteria</p> <p>1. Commercial Plans</p> <p>Mountain health Co-Op considers computer-assisted surgical navigation for orthopedic procedures is considered investigational.</p>

Section 1862(a)(1)(A) of the Social Security Act is the basis for denying payment for types of care, specific items, services, or procedures, not excluded by any other statutory clause, meeting all technical requirements for coverage, but are determined to be any of the following:

- Not generally accepted in the medical community as safe and effective in the setting and for the condition for which it is used,
- Not proven to be safe and effective based on peer review or scientific literature
- Experimental
- Not medically necessary in the particular case
- Furnished at a level, duration or frequency that is not medically appropriate
- Not furnished in accordance with accepted standards of medical practice, or

Not furnished in a setting appropriate to the patient's medical needs and condition.

Items and services must be established as safe and effective to be considered medically necessary.

That is, the items and services must be:

- Consistent with the symptoms or diagnosis of the illness or injury under treatment; and
- Necessary for, and consistent with, generally accepted professional medical standards of care (e.g., not experimental or investigational);and
- Not furnished primarily for the convenience of the patient, the provider or supplier; and
- Furnished at the most appropriate level that can be provided safely and effectively to the patient.

Medical devices that are not approved for marketing by the Food and Drug Administration (FDA) are considered investigational by Medicare and are not considered reasonable and necessary for the diagnosis or treatment of illness or injury, or to improve functioning of a malformed body member. Mountain Health Co-Op payments, therefore, may not be made for medical procedures and services performed using devices that have not been approved for marketing by the FDA or for those not included in an FDA-approved investigational (IDE) trial.

Background

Implant Alignment for Knee Arthroplasty

For total knee arthroplasty, malalignment is commonly defined as a variation of more than 3° from the targeted position. Proper implant alignment is believed to be an important factor for minimizing long-term wear, the risk of osteolysis, and loosening of the prosthesis.

Computer-Assisted Navigation

The goal of computer-assisted navigation is to increase surgical accuracy and reduce the

chance of malposition.

In addition to reducing the risk of substantial malalignment, computer-assisted navigation may improve soft tissue balance and patellar tracking. Computer-assisted navigation is also being investigated for surgical procedures with limited visibility such as placement of the acetabular cup in total hip arthroplasty, resection of pelvic tumors, and minimally invasive orthopedic procedures. Other potential uses of computer-assisted navigation for surgical procedures of the appendicular skeleton include screw placement for fixation of femoral neck fractures, high tibial osteotomy, and tunnel alignment during the reconstruction of the anterior cruciate ligament.

Computer-assisted navigation devices may be image-based or non-image-based. Image-based devices use preoperative computed tomography scans and operative fluoroscopy to direct implant positioning. Newer non-image-based devices use information obtained in the operating room, typically with infrared probes. For total knee arthroplasty, specific anatomic reference points are made by fixing signaling transducers with pins into the femur and tibia. Signal emitting cameras (e.g., infrared) detect the reflected signals and transmit the data to a dedicated computer. During the surgery, multiple surface points are taken from the distal femoral surfaces, tibial plateaus, and medial and lateral epicondyles. The femoral head center is typically calculated by kinematic methods that involve the movement of the thigh through a series of circular arcs, with the computer producing a 3-dimensional model that includes the mechanical, trans epicondylar, and tibial rotational axes. Computer-assisted navigation systems direct the positioning of the cutting blocks and placement of the prosthetic implants based on the digitized surface points and model of the bones in space. The accuracy of each step of the operation (cutting block placement, saw cut accuracy, seating of the implants) can be verified, thereby allowing adjustments to be made during surgery. For spine surgery, computer-assisted navigation may improve the accuracy of pedicle screw placement compared to conventional screw placement methods and limit radiation exposure to patients and surgical teams.

Computer-assisted navigation involves 3 steps: data acquisition, registration, and tracking.

Data Acquisition

Data can be acquired in 3 ways: fluoroscopically, guided by computed tomography scan or magnetic resonance imaging, or guided by imageless systems. These data are then used for registration and tracking.

Registration

Registration refers to the ability to relate images (i.e., radiographs, computed tomography scans, magnetic resonance imaging, or patients' 3-dimensional anatomy) to the anatomic position in the surgical field. Registration techniques may require the placement of pins or "fiducial markers" in the target bone. A surface-matching technique can also be used in which the shapes of the bone surface model generated from preoperative images are matched to surface data points collected during surgery.

Tracking

Tracking refers to the sensors and measurement devices that can provide feedback during surgery regarding the orientation and relative position of tools to bone anatomy. For example, optical or electromagnetic trackers can be attached to regular surgical tools, which then provide real-time information of the position and orientation of tool alignment concerning the bony anatomy of interest.

VERASENSE™ (OrthoSense) is a single-use device that replaces the standard plastic tibial trial spacer used in total knee arthroplasty. The device contains microprocessor sensors that quantify load and contact position of the femur on the tibia after resections have been made. The wireless sensors send the data to a graphic user interface that depicts the load. The device is intended to provide quantitative data on the alignment of the implant and soft tissue balancing in place of intraoperative "feel."

iASSIST® (Zimmer) is an accelerometer-based alignment system with a user interface built into disposable electronic pods that attach to the femoral and tibial alignment and resection guides. For the tibia, the alignment guide is fixed between the tibial spines and a claw on the malleoli. The relation between the electronic pod of the digitizer and the bone reference is registered by moving the limb into abduction, adduction, and neutral position. Once the information has been registered, the digitizer is removed, and the registration data are transferred to the electronic pod on the cutting guide. The cutting guide can be adjusted for varus/valgus alignment and tibial slope. A similar process is used for the femur. The pods use the wireless exchange of data and display the alignment information to the surgeon within the surgical field. A computer controller must also be present in the operating room.

Due to the lack of any recent studies on pelvic tumor resection, these sections of the Rationale were removed from this evidence review in 2016.

Regulatory Status

Because computer-assisted navigation is a surgical information system in which the surgeon is only acting on the information that is provided by the navigation system, surgical navigation systems generally are subject only to 510(k) clearances from the U.S. Food and Drug

Administration (FDA). As such, the FDA does not require data documenting the intermediate or final health outcomes associated with computer-assisted navigation. In contrast, robotic procedures, in which the actual surgery is robotically performed, are subject to the more rigorous requirement of the process.

A variety of surgical navigation procedures have been cleared for marketing by the FDA through the 510(k) process with broad labeled indications. For example, The OEC FluoroTrak 9800 plus is marketed for locating anatomic structures anywhere on the human body.

Several navigation systems (e.g., PiGalileo™ Computer-Assisted Orthopedic Surgery System, PLUS

Orthopedics; OrthoPilot® Navigation System, Braun; Navitrack® Navigation System, ORTHOsoft) have received the FDA clearance specifically for total knee arthroplasty. The FDA cleared indications for the PiGalileo system are representative. This system "is intended to be used in computer-assisted orthopedic surgery to aid the surgeon with bone cuts and implant positioning during joint replacement. It provides information to the surgeon that is used to place surgical instruments during surgery using anatomical landmarks and other data specifically obtained intraoperatively (e.g., ligament tension, limb alignment). Examples of some surgical procedures include but are not limited to:

- Total knee replacement supporting both bone referencing and ligament balancing techniques
- Minimally invasive total knee replacement."

FDA product code: HAW.

In 2013, the VERASENSE Knee System (OrthoSensor) and the iASSIST Knee (Zimmer) were cleared for marketing by the FDA through the 510(k) process. FDA product codes: ONN, OLO.

Several computer-assisted navigation devices cleared by the FDA are listed in the table below.

Table 1. Computer-Assisted Navigation Devices Cleared by the U.S. Food and Drug Administration

Device	Manufacturer	Date Cleared	510(k) No.	Indication
Vital™ Navigation System	Zimmer Biomet Spine, Inc.	12/02/2019	K191722	Computer-assisted Navigation for Orthopedic Surgery
Stryker Navigation System With Spinemap Go Software Application, Fluoroscopy Trackers And Fluoroscopy Adapters. Spinemask Tracker	Stryker Corporation	02/14/2019	K183196	Computer-assisted Navigation for Orthopedic Surgery
NuVasive Pulse™ System	NuVasive Inc.	6/29/2018	K180038	Computer-assisted Navigation for Orthopedic Surgery
VERASENSE for Zimmer Biomet Persona	OrthoSensor Inc.	6/7/2018	K180459	Computer-assisted Navigation for Orthopedic Surgery

StealthStation™ S8 With Spine Software	Medtronic	5/01/2017	K170011	Computer-assisted Navigation for Orthopedic Surgery
NuVasive Next Generation NVM5® System	NUVASIVE Inc.	3/16/2017	K162313	Computer-assisted Navigation for Orthopedic Surgery
Stryker OrthoMap Versatile Hip System	Stryker Corporation	2/23/2017	K162937	Computer-assisted Navigation for Orthopedic Surgery
JointPoint™	JointPoint Inc.	8/3/2016	K160284	Computer-assisted Navigation for Orthopedic Surgery
ExactechGPS®	Blue Ortho	7/13/2016	K152764	Computer-assisted Navigation for Orthopedic Surgery
Verasense Knee System	OrthoSensor Inc.	4/15/2016	K150372	Computer-assisted Navigation for Orthopedic Surgery
iASSIST Knee System	Zimmer CAS	9/11/2014	K141601	Computer-assisted Navigation for Orthopedic Surgery
CTC TCAT(R)-TPLAN(R) Surgical System	Curexo Technology Corporation	8/18/2014	K140585	Computer-assisted Navigation for Orthopedic Surgery
Digimatch™ Orthodoc Robodoc® Encore Surgical System	Curexo Technology Corporation	5/27/2014	K140038	Computer-assisted Navigation for Orthopedic Surgery

Rationale

This evidence review was created in February 2004 and has been updated regularly using with searches of the PubMed database. The most recent literature update was performed through February 13, 2024.

Evidence reviews assess clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

For many orthopedic surgical procedures, optimal alignment is considered an important aspect of long-term success. For example, misplaced tunnels in the anterior cruciate ligament or posterior cruciate ligament reconstruction or malalignment of arthroplasty components are some of the leading causes of instability and reoperation. In total hip arthroplasty (THA), the orientation of the acetabular component of the THA is considered critical, while for total knee arthroplasty (TKA), alignment of the femoral and tibial components and ligament balancing are considered important outcomes. Ideally, one would prefer controlled trials comparing the longterm outcomes, including stability and reoperation rates.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

Computer-Assisted Navigation for Trauma or Fracture

Clinical Context and Therapy Purpose

The purpose of computer-assisted navigation is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conventional/manual alignment methods, in individuals who are undergoing orthopedic surgery for trauma or fracture. The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing orthopedic surgery for trauma or fracture.

Interventions

The therapy being considered is computer-assisted navigation. Computer-assisted navigation in

orthopedic procedures describes the use of computer-enabled tracking systems to facilitate alignment in a variety of surgical procedures, including fixation of fractures, ligament reconstruction, osteotomy, tumor resection, preparation of the bone for joint arthroplasty, and verification of the intended implant placement.

Comparators

Comparators of interest include conventional/manual alignment methods. Treatment by means of conventional/manual alignment methods include medical reduction procedures and conventional fluoroscopic guidance (i.e., C-arm fluroscopy).

Outcomes

The general outcomes of interest are symptoms, morbid events, and functional outcomes.

The existing literature evaluating computer-assisted navigation as a treatment for patients who are undergoing orthopedic surgery for trauma or fracture has varying lengths of follow-up. While studies described below all reported at least one outcome of interest, longer follow-up was necessary to fully observe outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Computer-assisted surgery has been described as an adjunct to pelvic, acetabular, or femoral fractures. For example, fixation of these fractures typically requires percutaneous placement of screws or guidewires. Conventional fluoroscopic guidance (i.e., C-arm fluoroscopy) provides imaging in only 1 plane. Therefore, the surgeon must position the implant in 1 plane and then get additional images in other planes in a trial-and-error fashion to ensure that the device has been properly placed. This process adds significant time in the operating room and radiation exposure. Computer-assisted surgery may permit minimally invasive fixation and provide more versatile screw trajectories with less radiation exposure. Therefore, computed-assisted surgery is considered an alternative to the existing image guidance using C-arm fluoroscopy.

Observational Study

Ideally, investigators would conduct controlled trials comparing operating room time,

radiation exposure, and long-term outcomes of those whose surgery was conventionally guided using Carm versus image-guided using computer-assisted surgery. While several in vitro and review studies had been published,^{1,2,3} only 2 studies of computer-assisted surgery in trauma or fracture cases were identified.^{4,5} Computer-assisted navigation for internal fixation of femoral neck fractures was retrospectively analyzed in 2 cohorts of consecutive patients (20 each, performed from 2001 to 2003, at 2 different campuses of a medical center) who underwent internal fixation with 3 screws for a femoral neck fracture.⁴ Three of 5 measurements of parallelism and neck coverage were significantly improved by computer-assisted navigation; they included a larger relative neck area held by the screws (32% vs. 23%) and less deviation on the lateral projection for both the shaft (1.7° vs. 5.2°) and the fracture (1.7° vs. 5.5°) screw angles, all respectively. Slight improvements in anteroposterior screw angles (1.3° vs. 2.1° and 1.3° vs. 2.4°, respectively) were not statistically significant. There were 2 reoperations in the computer-assisted navigation group and 6 in the conventional group. Complications (collapse, subtrochanteric fracture, head penetration, osteonecrosis) were lower in the computer assisted navigation group (3 vs. 11, respectively).

A retrospective comparative study by Swartman et al (2021) investigated differences in conventional fluoroscopy-assisted percutaneous management (n=13) of acetabular fractures to 3-dimensional (3D) -computer navigated management (n=24).⁵ Both groups demonstrated significant reduction in fracture gaps and steps post-intervention. However, there were no significant differences between groups in outcomes related to fracture reduction or screw positions.

Section Summary: Computer-Assisted Navigation for Trauma or Fracture

There is limited literature on the use of computer-assisted navigation for trauma or fractures. Additional controlled studies that measure health outcomes are needed to evaluate this technology.

Computer-Assisted Navigation for Anterior Cruciate Ligament or Posterior Cruciate Ligament Reconstruction

Clinical Context and Therapy Purpose

The purpose of computer-assisted navigation is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conventional/manual alignment methods, in individuals who are undergoing ligament reconstruction.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing ligament reconstruction.

Interventions

The therapy being considered is computer-assisted navigation. Computer-assisted navigation in orthopedic procedures describes the use of computer-enabled tracking systems to facilitate alignment

in a variety of surgical procedures, including fixation of fractures, ligament reconstruction, osteotomy, tumor resection, preparation of the bone for joint arthroplasty, and verification of the intended implant placement.

Comparators

Comparators of interest include conventional/manual alignment methods. Treatment by means of conventional/manual alignment methods include medical reduction procedures.

Outcomes

The general outcomes of interest are symptoms, morbid events, and functional outcomes. The existing literature evaluating computer-assisted navigation as a treatment for patients who are undergoing ligament reconstruction has varying lengths of follow-up. While studies described below all reported at least 1 outcome of interest, longer follow-up was necessary to fully observe outcomes. Therefore, 2 years of follow-up is considered necessary to demonstrate efficacy.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

Eggerding et al (2014) published a Cochrane review that compared the effects of computer-assisted navigation with conventional operating techniques for anterior cruciate ligament or posterior cruciate ligament reconstruction.⁶ Five RCTs (N=366 participants) on anterior cruciate ligament reconstruction were included in the updated review; no studies involved posterior cruciate ligament reconstruction. The quality of evidence ranged from moderate to very-low. Pooled data showed no statistically or clinically relevant differences in self-reported health outcomes (International Knee Documentation Committee [IKDC] subjective scores and Lysholm Knee Scale scores) at 2 or more years of follow-up. No significant differences were found for secondary outcomes, including knee stability, range of motion, and tunnel placement. Overall, there was insufficient evidence to advise for or against the use of computer-assisted navigation. Four of the 5 trials included in the Cochrane review are described in Tables 2 and 3.

Yavari et al (2023) published a systematic review of 11 studies (N=775) evaluating technology-assisted anterior cruciate ligament reconstruction including computer-assisted navigation, virtual reality,

augmented reality, 3D printing, and robotics.⁷ Five studies (N=454) evaluated image-free computer-assisted navigation. Subjective IKDC scores were improved in the technology-assisted surgery group (mean difference, 1.97; 95% CI, 0.27 to 3.66; p=.02); however, the authors noted that a minimally clinically important difference of 9 has been previously published. Objective IKDC scores were not significantly different between groups (risk ratio, 1.02; 95% CI, 0.98 to 1.06). Notably, results specific to computer-assisted navigation were not analyzed.

Randomized Controlled Trials

Plaweski et al (2006) reported on a trial that randomized 60 patients to manual or computer-assisted guidance for tunnel placement with follow-up at 1, 3, 6, 12, 18, and 24 months.⁸ There were no differences between groups in measurements of laxity. However, there was less variability in side-to-side anterior laxity in the navigated group (e.g., 97% were within 2 mm of laxity in the navigated group vs. 83% in the conventional group at an applied force of 150 N). There was a significant difference in the sagittal position of the tibial tunnel (distance from the Blumensaat line, 0.4 mm vs. -1.2 mm, respectively), suggesting possible impingement in extension for the conventional group. At the final follow-up (24 months), all knees had normal function, with no differences observed between groups.

Hart et al (2008) compared biomechanical radiographic with functional results in 80 patients randomized to anterior cruciate ligament reconstruction using computer-assisted navigation (n=40) or to the standard manual targeting technique (n=40).⁹ The blinded evaluation found more exact bone tunnel placement with computer-assisted navigation, but no overall difference in biomechanical stability or function between groups.

Other studies have found no significant improvement in the accuracy of tunnel placement when using computer-assisted navigation. Meuffels et al (2012) reported on a double-blind controlled trial that randomized 100 patients to conventional or computer-assisted surgery.¹⁰ Evaluation by 3-dimensional computed tomography (CT) found no significant difference between groups for the accuracy or the precision of the femoral and tibial tunnel placement.

Table 2. Summary of Characteristics of Key Randomized Controlled Trials Comparing Computer-Assisted Navigation with Manual Placement for Anterior or Posterior Cruciate Ligament Reconstruction

Study	Countries	Sites	Dates	Participates	Interventions	

					Active	Comparator
Plaweski et al (2006) ^{8,}	USA	1	Oct 2014 to Jan 2016	Patients (N=60) undergoing ACL reconstruction.	CAN (n=30)	Manual placement (n=30)
Hart et al (2008) ^{9,}	Czech Republic	1	NR	Patients (N=80) undergoing ACL reconstruction for chronic rupture of the ACL; only chronic ACL-insufficiency knees were included in the study (>6 mo after injury). Other inclusion criteria were no other prior or simultaneous intra-articular surgical procedure, no cartilage degeneration of meniscal tear, and a normal contralateral knee. Ages ranged from 16 to 39 years with a mean of 29.4. Mean body weight was 74 kg.	CAN (n=40)	Manual placement (n=40)
Meuffels et al (2012) ^{10,}	Netherlands	1	Jan 2007 to Nov 2009	Patients (N =100) ≥18 years of age and eligible for primary ACL reconstruction without additional PCL or lateral collateral ligament injury were included.	CAN (n=49)	Conventional (n=51)

Mauch et al (2007) ¹¹ ,	Germany	1	Dec 2003 to April 2004	Athletes aged 18 to 49 years (N=53) with ACL rupture and no complex injuries of knee with additional injury of PCL, injury of posterior lateral complex, or third-degree injury of intra-articular ligament.	CAN (n=24)	Manual placement (n=29)
------------------------------------	---------	---	------------------------	--	------------	-------------------------

ACL: anterior cruciate ligament; CAN: computer-assisted navigation; NR: not reported; PCL: posterior cruciate ligament.

Table 3. Summary of Key Randomized Controlled Trials Comparing Computer-Assisted Navigation With Manual Placement for Anterior or Posterior Cruciate Ligament Reconstruction

Study	IKDC	Laxity <2 mm	Lachman Test (0)	Lachman Test (2+)	Placement of the Femoral Tunnel	Tibial Tunnel Border
Plaweski et al (2006) ⁸ ,						
CAN (n=26 knees)					NR	mean ATB, - 0.2 (5 to +4)
Mean Level A laxity level (n=26 knees)	mean, 1.3 mm at 200 N; p=.49	96.7%; p=.295	23 (76.7)	1 (3.3)		
Manual (n=22 knees)					NR	mean ATB, 0.4 (0 to 3)
Mean Level A laxity level (n=22 knees)	mean, 1.5 mm at 200 N; p=.49	83%; p=.292	26 (87)	0 (0)		
Hart et al (2008) ⁹ ,						

CAN (n=40)	Mean post-op Improvement: 76.5 points; SD, 10.3; p<.01	Mean difference in anterior laxity Compared with contralateral (healthy) knee: 1.43 mm (range, 0 to 4 mm)	12 (30%)	14 (35%)	Ideal a/t value: 24.8% Mean, 25.5% (SD, 1.63)	Zone 2 location: 39
Manual (n=40)	Mean post-op Improvement: 73.1 points; SD, 11.8; p<.01	Mean difference in anterior laxity compared with contralateral (healthy) knee: 1.24 mm (range, - 2 to 5 mm)	18 (45%)	10 (25%)	Ideal a/t value: 24.8% Mean, 27.% (SD, 2.76)	Zone 2 location: 38 (95.0%)
Meuffels et al (2012)10,						
CAN (n=49)	NR	NR	NR	NR	Mean 39% of the proximal distance on the intracondylar axis	Distance from most medial edge: 42.7% ±3.6%
Manual (n=51)	NR	NR	NR	NR	Mean 39.7% of the proximal distance on the intracondylar axis	Distance from most medial edge: 42.6% ±5.7%
Mauch et al (2007)11,						

CAN (n=24)	NR	NR	NR	NR	NR	21.2 mm (32.2%)
Manual (n=29)	NR	NR	NR	NR	NR	19.4 mm)

a/t value: ratio identifies anterior-posterior femoral tunnel placement; ATB: anterior tension band plate; CAN: computer-assisted navigation; IKDC: International Knee Documentation Committee; NR: not reported; Post-op: postoperative; SD: standard deviation.

The purpose of the limitations tables (see Tables 4 and 5) is to display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence following each table and provides the conclusions on the sufficiency of evidence supporting the position statement.

Table 4. Summary of Study Relevance Limitations in Key Randomized Controlled Trials Comparing Computer-Assisted Navigation With Manual Placement

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Plaweski et al (2006) ⁸ ,	3. Limited demographic information provided				
Hart et al (2008) ⁹ ,	3. the study setting and source of study participants are missing (as is the referral pattern) – this could create referral filter bias.				
Meuffels et al (2012) ¹⁰ ,	3. Study population is incompletely characterized.	2. Inconsistent fidelity of intervention protocol: There is a lack of consistency as to the best method for performing the			

		intervention.			
Mauch et al (2007) ¹¹ ,	1,4. Intended use population is unclear. Limited to athletes.			5,6. Clinically significant difference not prespecified or mentioned.	1,2. Follow-up was 4 days, not long enough to determine intermediate- or long-term outcomes.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other. b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest. Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not established and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported. e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 5. Summary of Design and Conduct Limitations in Key Randomized Controlled Trials Comparing Computer-Assisted Navigation with Manual Placement

	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Plaweski et al (2006) ⁸ ,		1. Unclear whether patients were blinded.				3. Confidence intervals not reported. 4. Comparison of treatment effect not provided.
Hart et al (2008) ⁹ ,	3. Randomization techniques are not described in any manner within the				1. Power calculations not reported.	3. Confidence intervals not reported.

	text.					
Meuffels et al (2012) ¹⁰ ,						
Mauch et al (2007) ¹¹ ,	4. Drawing lots is a weak method of allocation	1,2,3 Blinding is not mentioned at all			1. Power calculations not reported	3. Confidence intervals not reported

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4.

Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Section Summary: Computer-Assisted Navigation for Anterior Cruciate Ligament or Posterior Cruciate Ligament Reconstruction

The evidence on computer-assisted navigation for anterior cruciate ligament or posterior cruciate ligament reconstruction includes a systematic review of 5 RCTs. These RCTs, of moderate- to low-quality, did not consistently demonstrate more accurate tunnel placement with computer-assisted navigation. No studies have shown an improvement in functional outcomes or need for revision when computer-assisted navigation is used for anterior cruciate ligament or posterior cruciate ligament reconstruction.

Computer-Assisted Navigation for Total Hip Arthroplasty and Periacetabular Osteotomy Clinical Context and Therapy Purpose

The purpose of computer-assisted navigation is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conventional/manual alignment methods, in individuals who are undergoing THA and periacetabular osteotomy. The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing THA and periacetabular osteotomy.

Interventions

The therapy being considered is computer-assisted navigation. Computer-assisted navigation in orthopedic procedures describes the use of computer-enabled tracking systems to facilitate alignment in a variety of surgical procedures, including fixation of fractures, ligament reconstruction, osteotomy, tumor resection, preparation of the bone for joint arthroplasty, and verification of the intended implant placement.

Comparators

Comparators of interest include conventional/manual alignment methods. Treatment by means of conventional/manual alignment methods include medical reduction procedures.

Outcomes

The general outcomes of interest are symptoms, morbid events, and functional outcomes.

The existing literature evaluating computer-assisted navigation as a treatment for patients who are undergoing THA and periacetabular osteotomy has varying lengths of follow-up, ranging from 6 to 40 months. While studies described below all reported at least 1 outcome of interest, longer follow-up was necessary to fully observe outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

Kunze et al (2022) published a systematic review comparing surgical time, short-term adverse events, and implant placement accuracy between manual, robotic-assisted, and computernavigated THA.¹² Seven RCTs were identified comparing computer-assisted navigation and manual THAs. Table 6 outlines the studies included comparing computer-assisted THA and manual THA. Characteristics and results specific to computer-assisted navigation are shown in Tables 7 and 8, respectively. In brief, manual THA resulted in significantly shorter surgical times and a similar incidence of complications and revisions compared to computer-assisted THA. However, computer-assisted navigation THA led to increased precision in the placement of acetabular implants. These results are limited by a lack of recent RCTs, inability to conduct meta-analysis of patient-reported outcome measures, and use of the Lewinnek safe zone as a benchmark for proper acetabular implant positioning, which may not be appropriate in all patients. Additionally, there were a variety of computer-assisted navigation systems used across RCTs, limiting conclusions regarding any particular system.

Table 6. Comparison of RCTs Included in Systematic Reviews and Meta-Analyses for Computer-Assisted Navigation for THA

Study	Kunze et al (2022) ^{a12,}
Leenders et al (2002)	●
Gurgel et al (2014)	●
Lass et al (2014)	●
Renkawitz et al (2015)	●
Parratte et al (2016)	●
Weber et al (2016)	●
Verdier et al (2016(=)	●

RCT: randomized controlled trial; THA: total hip arthroplasty. a Only articles comparing computer navigation and manual total hip arthroplasty procedures are included in table.

Table 7. Systematic Reviews and Meta-Analyses for Computer-Assisted Navigation for THA: Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration, mean
-------	-------	--------	--------------	-----------	--------	----------------

Kunze et al (2022)¹²,	2008-2019	7	Computer-assisted navigation THA compared to manual THA with at least 1-year follow-up	598 (40 to 135)	RCT	4.3 yrs (range, 1 to 14.2 yrs) ^a
---	-----------	---	--	-----------------	-----	---

^a Mean duration includes all studies included in systematic review, including robotic-assisted THA studies. RCT: randomized controlled trial; THA: total hip arthroplasty

Table 8. Systematic Reviews and Meta-Analyses for Computer-Assisted Navigation for THA: Results

Study	Operation length, mins	All-cause complications	All-cause revisions	Acetabular implant positioning (% of acetabular cups placed in safe zone)
Kunze et al (2022)¹²,				
Total N	373 (3 studies)	NR (11 studies)	598 (7 studies)	178 (3 studies)
Manual THA	mean, 86.6 mins	Total=38 (6.6%)	Total=2	46/89 (52%)
Computer-assisted THA	mean, 95.7 mins	Total=5 (1.7%)	Total=3	70/89 (79%)
Pooled effect (95% CI)	SMD, 8.55 (3.49 to 13.60)	OR, 0.83 (0.23 to 2.99)	OR, 1.15 (0.30 to 4.42)	ES, 0.79 (0.69 to 0.86)
p-value	<.001	.781	.840	.02
I ² (p)	--	--	--	0% (.29)

CI: confidence interval; ES: effect size; NR: not reported; OR: odds ratio; SMD: standard mean difference; THA: total hip arthroplasty.

Nonrandomized Studies

Manzotti et al (2011) compared leg length restoration in a matched-pair study.¹³ Forty-eight patients undergoing THA with computer-assisted navigation were compared with patients who were matched for age, sex, arthritis level, preoperative diagnosis, and preoperative leg length discrepancy and underwent conventional freehand THA using the same implant in the same period. The mean preoperative leg length discrepancy was 12.17 mm in the computer-assisted navigation group and 11.94 mm in the standard group. Surgical time was increased by 16 minutes in the computer-assisted navigation group (89 minutes vs. 73 minutes). There was a significant decrease in both the mean postoperative leg length discrepancy (5.06 mm vs. 7.65 mm) and the number of cases with a leg length discrepancy of 10 mm or more (5 patients vs. 13 patients), all respectively.

Outcomes at 40-month follow-up (range, 7 to 77 months) did not differ significantly for the Harris Hip Score (88.87 vs. 89.73) or the 100-point normalized Western Ontario and McMaster Universities Arthritis Index score (9.33 vs. 13.21; $p=.050$), all respectively. Longer follow-up with a larger number of subjects is needed to determine whether computer-assisted navigation influences clinical outcomes.

Minimally Invasive Total Hip Arthroplasty

Systematic Reviews

It has been proposed that computer-assisted navigation might overcome the difficulties of reduced visibility of the surgical area associated with minimally invasive procedures. Ulrich et al (2007) summarized study results that compared outcomes from minimally invasive THA using computer-assisted navigation with standard THA.¹⁴ Seventeen studies were described in this evidence-based review, including 9 prospective comparisons, 7 retrospective comparisons, and 1 large (N=100) case series. Reviewers concluded that alignment with minimally invasive computer-assisted navigation appears to be at least as good as standard THA, although the more consistent alignment must be balanced against the expense of the computer systems and increased surgical time.

Randomized Controlled Trials

Reininga et al (2013) reported short-term outcomes of minimally invasive THA approach with computer-assisted navigation (n=35) compared with conventional posterolateral THA (n=40).¹⁵ This randomized comparison found no group differences in the recovery of gait at up to 6 months post-surgery.

Periacetabular Osteotomy

Randomized Controlled Trials

Hsieh et al (2006) reported on 36 patients with symptomatic adult dysplastic hip who were randomized to CT-based navigation or the conventional technique for periacetabular osteotomy.¹⁶ An average of 0.6 intraoperative radiographs were taken in the navigated group compared with 4.4 in the conventional group, resulting in a total surgical time that was 21 minutes shorter for computer-assisted navigation. There were no differences between groups for correction in femoral head coverage or functional outcomes (pain, walking, range of motion) at 24 months.

Total Hip Resurfacing

Randomized Controlled Trials

Stiehler et al (2013) reported on short-term radiographic and functional outcomes from a randomized comparative trial of total hip resurfacing using computer-assisted navigation and conventional total hip resurfacing in 75 patients.¹⁷ For most of the radiographic measures, there were no significant differences between the computer-assisted navigation and conventional total hip resurfacing groups. There were fewer outliers ($\geq 5^\circ$) for the femoral component with computer-assisted navigation (11%)

compared with conventional placement (32%). At 6-month follow-up, there were no differences between groups in the final Western Ontario and McMaster Universities score or Harris Hip Score. The computer-assisted navigation group did show a greater percentage improvement in the Western Ontario and McMaster Universities scores and Harris Hip Score due to differences between groups at baseline.

Section Summary: Computer-Assisted Navigation for Total Hip Arthroplasty and Periacetabular Osteotomy

Relatively few RCTs have evaluated computer-assisted navigation for hip procedures. Although there was an early interest in this technology, no recent RCTs have been identified. There is inconsistent evidence from systematic reviews of these small trials on whether computer assisted navigation improves alignment with conventional or minimally invasive THA. One RCT found improved alignment when computer-assisted navigation was used for hip resurfacing, but there was little evidence of improved outcomes at short-term follow-up. Overall, improved health outcomes have not been demonstrated with computer-assisted navigation for any hip procedures.

Computer-Assisted Navigation for Total Knee Arthroplasty

Clinical Context and Therapy Purpose

The purpose of computer-assisted navigation is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conventional/manual alignment methods, in individuals who are undergoing TKA.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing TKA.

Interventions

The therapy being considered is computer-assisted navigation. Computer-assisted navigation in orthopedic procedures describes the use of computer-enabled tracking systems to facilitate alignment in a variety of surgical procedures, including fixation of fractures, ligament reconstruction, osteotomy, tumor resection, preparation of the bone for joint arthroplasty, and verification of the intended implant placement.

Comparators

Comparators of interest include conventional/manual alignment methods. Treatment by means of conventional/manual alignment methods include medical reduction procedures.

Outcomes

The general outcomes of interest are symptoms, morbid events, and functional

outcomes. The existing literature evaluating computer-assisted navigation as a treatment for patients who are undergoing TKA has varying lengths of follow-up, ranging from 1 to 8 years. While studies described below all reported at least 1 outcome of interest, longer follow-up was necessary to fully observe outcomes.

Alignment of a knee prosthesis can be measured along several different axes, including the mechanical axis, and the frontal and sagittal axes of both the femur and tibia.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

A systematic review conducted by Xie et al (2012) included 21 randomized trials (N=2658 patients) that reported on clinical outcomes with or without the use of computer-assisted navigation (Table 9).¹⁸ Most trials included in the review had short-term follow-up. Surgical time was significantly increased with computer-assisted navigation for TKA, but there was no significant difference between approaches in total operative blood loss, the Knee Society Score, or range of motion (Table 10).

Rebal et al (2014) conducted a meta-analysis of 20 RCTs (N=1713 knees) that compared imageless navigation technology with conventional manual guides (Table 9).¹⁹ The majority of included studies had a low risk of bias. The improvement in Knee Society Score was statistically superior in the computer-assisted navigation group at 3 months and 12 to 32 months (Table 10). However, these improvements did not achieve the minimal clinically significant difference, defined as a change of 34.5 points.

Table 9. Characteristics of Systematic Reviews and Meta-Analyses Investigating Total Knee Arthroplasty

Study	Dates	Trials	Participants	N (Range)	Design	Duration
-------	-------	--------	--------------	-----------	--------	----------

Xie et al (2012) ¹⁸ ,	PubMed and EMBASE through August 2011	21	Included 2658 patients. Among these, 1376 were randomly allocated to the computer-assisted TKA group and 1282 to the conventional group	2658 (25 to 120)	RCT	NR
Rebal et al (2014) ¹⁹ ,	PubMed, EMBASE, Scopus, and CENTRAL through December 2012	20	Included a combined 869 knees in the computer-assisted groups, and 844 knees in the control groups for a total of 1713 knees analyzed	1713 knees (46 to 166)	RCT	3 mos and 12 to 32 mos

NR: not reported; RCT: randomized controlled trial; TKA: total knee arthroplasty.

Table 10. Results of Systematic Reviews and Meta-Analyses Investigating Total Knee Arthroplasty

Study	Knee Society Score				Operative Time	
	CAN		Conventional		CAN (min)	Conventional (min)
Xie et al (2012) ¹⁸ ,						
Mean standard difference	4.47				14.68	
95% CI	-1.05 to 9.99				11.74 to 17.62	
P-value	.36				<.0001	
	CAN		Conventional		CAN (min)	Conventional (min)
	3 Months	12 to 32 Months	3 Months	12 to 32 Months		
Rebal et al (2014) ¹⁹ ,						
Mean	68.5	53.1	58.1	45.8	101.6	83.3
95% CI			1.13 to 19.78	2.87 to 11.90		11.84 to 24.60
P-value			.03	<.01		<.01

CAN: computer-assisted navigation; CI: confidence interval.

Effect of Computer-Assisted Navigation on Mid- to Long-Term Outcomes

Randomized Controlled Trials

RCTs comparing outcomes at 4 to 12 years follow-up generally have shown a reduction in the number of outliers with computer-assisted navigation, but little to no functional

difference between the computer-assisted navigation and conventional TKA groups.

Three trials comparing computer-assisted navigation and conventional surgery reported on outcomes at 4 to 5 years follow-up (N=67 to 107). Blakeney et al (2014) reporting 46-month follow-up for 107 patients, found a trend toward higher scores on the Oxford Knee Questionnaire with computer-assisted navigation, with a mean score of 40.6 for the computer assisted navigation group compared with 37.6 and 36.8 in extramedullary and intramedullary control groups, respectively. There were no significant differences in the 12-Item Short-Form Health Survey Physical Component or Mental Component Summary scores. The trial was underpowered, and the clinical significance of this trend for the Oxford Knee Questionnaire is unclear. Lutzner et al (2013), reporting on 5-year follow-up for 67 of 80 patients²¹, found a significant decrease in the number of outliers with computer-assisted navigation (3 vs. 9; $p=.048$) but no significant differences between groups on the Knee Society Score or Euroquol quality of life questionnaire. At 10-years post-surgery, a follow-up study (Beyer et al 2021) of 50 patients originally included in the Lutzner et al 2013 study showed no significant differences in the number of outliers between groups, patient-reported outcomes from the Knee Society Score of Euroquol quality of life questionnaire, and no differences in revision risk.²² Cip et al (2014) found a significant decrease in malalignment with computer-assisted navigation, but no significant differences in implant survival or consistent differences in clinical outcome measures between the navigated (n=100) and conventional (n=100) total knee arthroplasty groups at minimum 5-year follow-up.²³,

Four additional trials comparing computer-assisted and conventional surgery reported outcomes after 8 to 12 years follow-up (N=60 to 200). Hsu et al (2019) reported similar clinical and functional outcomes with the 2 procedures after a mean 8.1-year follow-up, although computer-assisted navigation achieved better radiographic alignment and fewer outliers.²⁴ They suggested that TKA with computer-assisted navigation may not provide an advantage to the typical osteoarthritis patient, but it may benefit certain patients, such as those with severe deformity of the knee joint, extra-articular deformities, and severe femoral bowing. The study was limited by its solely Asian patient population, single-center, and small sample size. Song et al (2016) also reported on a reduction in the number of outliers with computer-assisted navigation (7.3% vs. 20%; $p=.006$), with no significant differences in clinical outcomes at 8-year follow-up.²⁵ The trial, which assessed 80 patients (88 knees) was powered to detect a 3-point difference in Knee Society Score results. Cip et al (2018) published the results of a prospective randomized trial (N=200) comparing conventional TKA with computer-assisted TKA with a mean follow-up of 12 years postoperatively.²⁶ The trial was aimed at determining the long-term outcomes of computer-assisted navigation for TKA as a tool to expedite long-term survival based on improved postoperative implantation. The follow-up rate was 75%. No difference in long-term TKA survival was found between the conventional group (91.5%) and the computer assisted navigation group (98.2%) at 12 years ($p=.181$). In a single-blinded, prospective RCT, Farhan-Alanie et al (2023) compared conventional TKA (n=98) with computer-assisted TKA (n=101), with a mean follow-up of 10 years.²⁷ Over the 10-year period, there were 23 deaths (22.8%) in the computer-assisted group and 30 deaths (30.6%) in the conventional cohort. At the 10-year follow-up, the authors found

no difference in revision rates (4.0% computer navigation vs 6.1% conventional; $p=.429$) or clinical outcomes, including Oxford Knee Scores, American Knee Society Scores, or mental and physical scores on the 36-item Short-Form survey between groups.

Comparative Studies

Results from observational studies have generally been consistent with the systematic reviews and RCTs.^{28,29,30,31,32,33} The longest of these observational studies, conducted by Dyrhovden et al (2016), assessed survivorship and the relative risk of revision at 8-year follow-up for 23,684 cases from the Norwegian Arthroplasty Register for patients treated with computer-assisted navigation or conventional surgery.³² Overall prosthesis survival and risk of revision were similar for both groups, although revisions due to malalignment were reduced with computer-assisted navigation (relative risk, 0.5; 95% confidence interval [CI], 0.3 to 0.9; $p=.02$). There were no significant differences between groups for other reasons for revision (e.g., aseptic loosening, instability, periprosthetic fracture, decreased range of motion). At 8 years, the survival rate was 94.8% (95% CI, 93.8% to 95.8%) in the computer-assisted navigation group and 94.9% (95% CI, 94.5% to 95.3%) for conventional surgery.

In the largest observational study, Antonios et al (2020) compared Medicare data from 75,709 patients who underwent a computer navigated TKA with a matched cohort of 75,676 Medicare patients who underwent conventional TKA.³³ There was no statistically significant difference in 5-year event-free survival in all-cause revisions between groups (95.1% vs. 94.7%; $p=.06$) However, there was a small difference in revisions due to mechanical complications (96.1% vs. 95.7%; $p=.02$) but not in revisions due to periprosthetic joint infection (97.9% vs. 97.9%; $p=.30$)

A retrospective comparison cohort study by Webb et al (2021) compared conventional TKA cases ($n=219,880$) to computer navigated TKA cases ($n=5243$) that occurred from 2008 through 2016 and were documented in the American College of Surgeons National Surgical Quality Improvement Program database.³⁴ In univariate analysis of unmatched cohorts, rates of composite serious morbidities and death or serious morbidity were significantly higher in the conventional TKA group than the computer navigated group (8.47% vs. 7.54%; $p=.016$). In multivariable regression analysis, computer navigated TKA was found to be significantly associated with lower rates of serious morbidity (odds ratio [OR], 0.83; $p=.001$), death or serious morbidity (OR, 0.82; $p<.001$) and length of stay (OR, 0.86; $p=.024$). Propensity score matching identified 4811 case pairs of conventional versus computer navigated TKA. Propensity-matched analyses demonstrated no significant difference in mortality, length of operation time, length of stay, or rates of reoperation or readmission. The composite rate of complications was 18% less in the computer navigated group compared to the conventional TKA group ($p=.009$).

Section Summary: Computer-Assisted Navigation for Total Knee Arthroplasty

Based on systematic reviews, a large number of RCTs have assessed outcomes for TKA

using computer-assisted navigation or conventional TKA without computer-assisted navigation. Results are consistent in showing reductions in the proportion of outliers greater than 3° in alignment. Results from individual RCTs and cohort studies up to 12 years postoperatively have not shown that these differences in alignment lead to improved patient outcomes.

Computer-Assisted Navigation for Spine Surgery

Clinical Context and Therapy Purpose

The purpose of computer-assisted navigation is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conventional/manual methods, in individuals who are undergoing spine surgery, particularly spinal fusion surgery for radiculopathy and correction of spinal deformities (e.g. scoliosis). Spinal fusion may include the use of pedicle screws. Pedicle screws are a type of bone screw that, along with rods, is used to secure the vertebrae in a fixed position following fusion. Pedicle screws may be removed once healing has occurred, or they can be left in place. Pedicle screw placement accuracy is critical, as misplacement can cause a variety of complications, including pain and weakness or perforation leading to damage to surrounding nerves, soft tissues and bones.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing spine surgery. This can include patients undergoing cervical, thoracic or lumbar pedicle screw placement in association with spinal fusion surgery, due to trauma or for correction of spinal deformities, or patients undergoing spinal tumor resection.

Interventions

The therapy being considered is computer-assisted navigation. Computer-assisted navigation in orthopedic procedures describes the use of computer-enabled tracking systems to facilitate alignment in a variety of surgical procedures, including tumor resection and pedicle screw placement.

Comparators

Comparators of interest include conventional/manual surgical methods, such as fluoroscopically-guided freehand surgery.

Outcomes

The general outcomes of interest are symptoms, morbid events, and functional outcomes. Many studies report pedicle screw perforation (breach or encroachment into surrounding tissues, bones or organs) as a measure of procedural success. However, because not all screw perforations lead to symptoms or morbid events, revision surgeries would be a more relevant measure of clinical

outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Pedicle Screw Insertion for Spinal Fusion or Deformity Correction

Systematic Reviews

Numerous systematic reviews of mostly retrospective observational studies have assessed pedicle screw placement using computer-assisted navigation; however, evidence on health outcomes from the reviews is limited.^{35,36,37,38} In a 2018 review conducted by Staartjes et al, comparing computer-assisted navigation (n=1779 pedicle screws) and freehand placement (n=1809 pedicle screws) and the need for intraoperative revision, there was a nonsignificant trend favoring freehand placement based on an imprecise risk estimate (OR, 1.46 ; 95% CI, 0.30 to 7.17; I²=88%).³⁷ The same review found the need for postoperative revision was significantly lower with computer-assisted navigation versus freehand placement (OR, 0.31; 95% CI, 0.21 to 0.46; I²=0%). Another review, conducted by Perdomo-Pantoja et al (2019)³⁸, reported similar rates of screw placement accuracy with computer-assisted navigation (95.5%) and other placement methods (90.5% to 93.1%). Consistent with the RCT evidence discussed below, an older review by Shin et al (2012)³⁶ found a lower risk of pedicle screw perforation with computer-assisted navigation (6%; 287/4814) versus conventional, non-navigated screw placement (15%; 556/3725; risk ratio [RR], 0.39; 95% CI, 0.31 to 0.49; I²=49%). The review found no difference between navigated and non-navigated screw placement on operative time (-3.06 minutes; 95% CI, -35.60 to 29.48), estimated blood loss (-91.6 mL ; 95% CI, -185.95 to 3.24), or overall revision rate per screw insertion (1.44% vs. 2.03%; p=.11).

Randomized Controlled Trials

Three RCTs have compared pedicle screw insertion by computer-assisted navigation with conventional surgical techniques (Table 11). None of the trials reported health outcomes or post-surgical follow-up (Table 12). In the largest RCT, conducted by Laine et al (2000),³⁹ computer-assisted navigation was associated with longer surgical time than conventional surgery and fewer instances of pedicle screw perforation. A second, smaller RCT conducted by Rajasekaran et al (2007)⁴⁰ found pedicle screw placement using computer-assisted navigation associated with shorter placement time and a lower

rate of pedicle perforation relative to fluoroscopically guided placement. The third trial (N=21) compared the risk of patient and surgical team radiation exposure with pedicle screw placement using computer-assisted navigation with freehand, fluoroscopically-guided screw placement.⁴¹ The trial found significantly higher radiation exposure to the surgical team during freehand screw insertion (p<.01) with no difference between intervention groups and cumulative patient radiation dose. Tables 13 and 14 summarize key study relevance and design and conduct limitations.

Table 11. Summary of Key Randomized Controlled Trial Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Laine et al (2000) ³⁹ ,	Finland	1	1998- 1999	Patients undergoing thoracolumbar or lumbosacral fusion	Pedicle screw placement using computer-assisted navigation; n=41	Conventional pedicle screw placement; n=50
Rajasekaran et al (2007) ⁴⁰ ,	India	1	Not reported	Patients with scoliosis (40° to 80°) or kyphosis (≤90°) undergoing spinal deformity correction of the thoracic spine	Pedicle screw placement using computer-assisted navigation; n=17	Pedicle screw placement using fluoroscopic guidance; n=16
Villard et al (2014) ⁴¹ ,	Germany	1	Not reported	Patients undergoing lower thoracic and lumbar posterior transforaminal interbody fusion	Pedicle screw placement using computer-assisted navigation; n=10	Pedicle screw placement using fluoroscopic guidance as needed; n=11

Table 12. Summary of Key Randomized Controlled Trial Results

Study; Trial	Mean Insertion Time	Pedicle Screw Perforation	Radiation Exposure
Laine et al (2000) ³⁹ ,	n=91 (496 screws)	n=91 (496 screws)	--
Computer-assisted navigation	40.0 (SD 16) minutes total insertion time	4.6% (10/219)	Not reported
Conventional placement	28.7 (SD 17) minutes total insertion time	13.4% (37/277)	Not reported
p value	p=.001	p=.006	--
Rajasekaran et al (2007) ⁴⁰ ,	n=33 (478 screws)	n=33 (478 screws)	--
Computer-assisted navigation	2.4 (SD 0.7) minutes per screw	2.1% (5/242)	Not reported
Conventional placement	4.6 (S.D 1.1) minutes per screw	22.9% (54/236)	Not reported
P value	P<.001	P<.001	--
Computer-assisted navigation	Not reported	Not reported	888 (SD 449) cGy•cm ²
Conventional placement	Not reported	Not reported	1884 (SD 881) cGy•cm ²
P value	--	--	P=.73

SD: standard deviation.

Table 13. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Laine et al (2000) ³⁹ ,				1,2. The study reported on pedicle screw perforation but not clinical outcomes.	1,2. Follow-up was insufficient to assess benefits and harms

Rajasekaran et al (2007) ⁴⁰ ,				1,2. The study reported on pedicle screw perforation but not clinical outcomes.	1,2. Follow-up was insufficient to assess benefits and harms
Villard et al (2014) ⁴¹ ,				1,2. The study reported on radiation exposure but not clinical outcomes.	1,2. Follow-up was insufficient to assess benefits and harms

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest. c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported. e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 14. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Laine et al (2000) ³⁹ ,					1. Power calculations not reported	
Rajasekaran et al (2007) ⁴⁰ ,					1. Power calculations not reported	
Villard et al (2014) ⁴¹ ,	3. Allocation concealment is unclear	1,2. Not blinded to treatment				

		assignment or outcome assessment				
--	--	--	--	--	--	--

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias. b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference. f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Other Indications

The use of computer-assisted navigation for the treatment of spinal tumors has been reported in uncontrolled case series and case reports.^{42,43,44} Although the use of computer-assisted navigation appears safe for tumor resection based on these reports, evidence is too limited to draw any conclusions regarding the effect of computer-assisted navigation on health outcomes.

Section Summary - Computer-Assisted Navigation for Spine Surgery

Evidence from RCTs and larger observational studies found that computer-assisted navigation was associated with lower rates of pedicle screw perforation compared with other surgical placement methods. Evidence on clinical outcomes, including long-term health outcomes, is lacking.

Summary of Evidence

For individuals who are undergoing orthopedic surgery for trauma or fracture and receive computer-assisted navigation, the evidence includes 2 retrospective studies, reviews, and in vitro studies. Relevant outcomes are symptoms, morbid events, and functional outcomes. Functional outcomes were not included in the first clinical trial, although it did note fewer complications with computer-assisted navigation versus conventional methods. The second trial found no differences between groups in rates of fracture reduction or screw positions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing ligament reconstruction and receive computer-assisted navigation, the evidence includes a systematic review of 5 randomized controlled trials (RCTs) of computer-assisted navigation versus conventional surgery for anterior and posterior cruciate ligament. Relevant outcomes are symptoms, morbid events, and functional outcomes. Trial results showed no consistent improvement of

tunnel placement with computer-assisted navigation, and no trials looked at functional outcomes or need for revision surgery with computer-assisted navigation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing hip arthroplasty and periacetabular osteotomy and receive computer-assisted navigation, the evidence includes systematic reviews of older RCTs and comparison studies. Relevant outcomes are symptoms, morbid events, and functional outcomes. Evidence on the relative benefits of computer-assisted navigation with conventional or minimally invasive total hip arthroplasty (THA) is inconsistent, and more recent RCTs are lacking. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing total knee arthroplasty (TKA) and receive computer-assisted navigation, the evidence includes RCTs, systematic reviews of RCTs, and comparative studies. Relevant outcomes are symptoms, morbid events, and functional outcomes. The main difference found between TKA with computer-assisted navigation and TKA without computer assisted navigation is increased surgical time with computer-assisted navigation. Few differences in clinical and functional outcomes were seen at up to 12 years post-procedure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing spine surgery and receive computer-assisted navigation, the evidence includes RCTs, comparative observational studies, and systematic reviews of those observational studies. Relevant outcomes are symptoms, morbid events, and functional outcomes. Computer-assisted navigation for pedicle screw insertion was consistently associated with lower rates of screw perforation relative to other screw insertion methods, but evidence on clinical outcomes such as revision rate is inconsistent or lacking, including long term outcome follow-up. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2011 Input

In response to requests, input was received from 3 academic medical centers while this policy was under review in 2011. Input was mixed on whether computer-assisted navigation is considered

investigational. One reviewer provided additional references on high tibial osteotomy and pelvic tumor resection.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in ‘Supplemental Information’ if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Orthopedic Surgeons

The American Academy of Orthopedic Surgeons updated guidelines in 2022 on surgical management of osteoarthritis of the knee.⁴⁵ Related to computer-assisted surgical navigation, the guidelines state there is no difference in outcomes, function, or pain between computer navigation and conventional techniques for total knee arthroplasty (strength of evidence: strong; strength of recommendation: moderate), and make no specific recommendation related to its use. The guidelines note that the advantages of surgical navigation remain unclear.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 15.

Table 15. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT06062615	Randomized Pilot Study Investigating Early Functional Outcomes With the Use of Robotic Assisted Versus Conventional Total Knee Arthroplasty	30	Dec 2023
NCT06036212	A Prospective, Single-blind, Multi-Centre, Randomised Controlled Study to Evaluate the Clinical and Patient Reported Outcomes Following Unicompartmental Knee Arthroplasty With a Robotic Assisted Technique	280	Mar 2036

NCT03628378	Randomized, Controlled, Single Center Observational Study to Compare the Safety and Performance of Navigation-assisted OrthoPilot® Elite and Robotic-assisted MAKO® Total Knee Arthroplasty	140	Jun 2025
NCT03628378	Outcomes in Free-hand Versus Sensor-guided Balancing in Total Knee Arthroplasty: a Randomized Controlled Trial	130	Dec 2022
NCT02717299 ^a	Making Sense Out of Total Knee Sensor Assisted Technology: A Randomized Control Trial	78	Apr 2021 (recruitment status unknown)
NCT04960345	Comparison of Accuracy and Clinical Outcomes Between Brainlab Knee 3 Computer-assisted Navigation Systems and Conventional Instruments in TKA: a Prospective Cohort Study	188	Dec 2023
Unpublished			
NCT01469299 ^a	Prospective Study Measuring Clinical Outcomes of Knee Arthroplasty Using the VERASENSE™ Knee System	285	Dec 2016 (updated 01/11/17)
NCT03668756	Comparison of Computer-Assisted Navigation and Conventional Instrumentation for Bilateral Total Knee Arthroplasty: The Functional Outcome of Mid-Term Follow-up Study	56	Aug 2018
NCT02190435 ^a	Computer-Assisted Navigation for Intramedullary Nail Fixation of Intertrochanteric Femur Fractures	65	Jan 2016
NCT03817632 ^a	Prospective, Multicenter, Observational, Comparative Clinical Trial on the Equivalence of Two Different OrthoPilot® Navigation System Generations Applied for Computer-assisted Total Knee Arthroplasty	217	Oct 2022

NCT: national clinical trial. a Denotes industry-sponsored or cosponsored trial.

Essential Health Benefits

The Affordable Care Act (ACA) requires fully insured non-grandfathered individual and small group benefit plans to provide coverage for ten categories of Essential Health Benefits (“EHBs”), whether the benefit plans are offered through an Exchange or not. States can define EHBs for their respective state. States vary on how they define the term small group. In Idaho, a small group employer is defined as an employer with at least two but no more than fifty eligible employees on the first day of the plan or contract year, the majority of whom are employed in Idaho. Large group employers, whether they are self-funded or fully insured, are not required to offer EHBs, but may voluntarily offer them. The ACA requires any benefit plan offering EHBs to remove all dollar limits for EHBs.

Applicable Coding

The following codes and coding guidance are provided for general reference purposes only and may not be all-inclusive. The inclusion of a code does not guarantee or imply any right to member coverage or provider reimbursement, nor does its exclusion represent or imply that coverage or reimbursement is unavailable. All benefit coverage determinations are subject to the member-specific benefit plan documentation as well as additional terms and conditions, including but not limited to the written coverage position set forth in this medical policy, legal requirements, and other policies and guidelines, as applicable.

CPT	20985	Computer-assisted surgical navigational procedure for musculoskeletal procedures; image-less (List separately in addition to code for primary procedure)
	0054T	Computer-assisted musculoskeletal surgical navigational orthopedic procedure with image guidance based on fluoroscopic images (List separately in addition to code for primary procedure)
	0055T	Computer-assisted musculoskeletal surgical navigational orthopedic procedure with image guidance based on CT/MRI images (List separately in addition to code for primary procedure)
	61783	stereotactic computer-assisted (navigational) procedure; spinal (List separately in addition to code for primary procedure)
ICD-10-CM		Investigational for all codes
ICD-10-PCS	8E09XBZ	Computer Assisted Procedure of Head and Neck Region
	8E0WXBZ	Computer Assisted Procedure of Upper Extremity
	8E0YXBZ	Computer Assisted Procedure of Lower Extremity
Type of Service	Surgical	
Place of Service	Inpatient Outpatient	

Vendors

- **MedCom**
- **Health Plan Services**

References

1. Hofstetter R, Slomczykowski M, Krettek C, et al. Computer-assisted fluoroscopy-based reduction of femoral fractures and antetorsion correction. *Comput Aided Surg.* 2000; 5(5): 311-25. PMID 11169877
2. Schep NW, Broeders IA, van der Werken C. Computer assisted orthopaedic and trauma surgery. State of the art and future perspectives. *Injury.* May 2003; 34(4): 299-306. PMID 12667784
3. Slomczykowski MA, Hofstetter R, Sati M, et al. Novel computer-assisted fluoroscopy system for intraoperative guidance: feasibility study for distal locking of femoral nails. *J Orthop Trauma.* Feb 2001; 15(2): 122-31. PMID 11232651
4. Liebergall M, Ben-David D, Weil Y, et al. Computerized navigation for the internal fixation of femoral neck fractures. *J Bone Joint Surg Am.* Aug 2006; 88(8): 1748-54. PMID 16882897
5. Swartman B, Pelzer J, Beisemann N, et al. Fracture reduction and screw position after 3D-navigated and conventional fluoroscopy-assisted percutaneous management of acetabular fractures: a retrospective comparative study. *Arch Orthop Trauma Surg.* Apr 2021; 141(4): 593-602. PMID 32519074
6. Eggerding V, Reijman M, Scholten RJ, et al. Computer-assisted surgery for knee ligament reconstruction. *Cochrane Database Syst Rev.* Aug 04 2014; (8): CD007601. PMID 25088229
7. Yavari E, Moosa S, Cohen D, et al. Technology-assisted anterior cruciate ligament reconstruction improves tunnel placement but leads to no change in clinical outcomes: a systematic review and meta-analysis. *Knee Surg Sports Traumatol Arthrosc.* Oct 2023; 31(10): 4299-4311. PMID 37329370
8. Plaweski S, Cazal J, Rosell P, et al. Anterior cruciate ligament reconstruction using navigation: a comparative study on 60 patients. *Am J Sports Med.* Apr 2006; 34(4): 542-52. PMID 16556753
9. Hart R, Krejzla J, Sváb P, et al. Outcomes after conventional versus computer-navigated anterior cruciate ligament reconstruction. *Arthroscopy.* May 2008; 24(5): 569-78. PMID 18442690
10. Meuffels DE, Reijman M, Verhaar JA. Computer-assisted surgery is not more accurate or precise than conventional arthroscopic ACL reconstruction: a prospective randomized clinical trial. *J Bone Joint Surg Am.* Sep 05 2012; 94(17): 1538-45. PMID 22832975
11. Mauch F, Apic G, Becker U, et al. Differences in the placement of the tibial tunnel during reconstruction of the anterior cruciate ligament with and without computer-assisted navigation. *Am J Sports Med.* Nov 2007; 35(11): 1824-32. PMID 17878429
12. Kunze KN, Bovonratwet P, Polce EM, et al. Comparison of Surgical Time, Short-term Adverse Events, and Implant Placement Accuracy Between Manual, Robotic-assisted, and Computer-navigated Total Hip Arthroplasty: A Network Meta-analysis of Randomized Controlled Trials. *J Am Acad Orthop Surg Glob Res Rev.* Apr 01 2022; 6(4). PMID 35472191
13. Manzotti A, Cerveri P, De Momi E, et al. Does computer-assisted surgery benefit leg length restoration in total hip replacement? Navigation versus conventional freehand. *Int Orthop.* Jan 2011; 35(1): 19-24. PMID 19904533
14. Ulrich SD, Bonutti PM, Seyler TM, et al. Outcomes-based evaluations supporting computer-assisted surgery and minimally invasive surgery for total hip arthroplasty. *Expert Rev Med Devices.* Nov 2007; 4(6): 873-83. PMID 18035952
15. Reininga IH, Stevens M, Wagenmakers R, et al. Comparison of gait in patients following a computer-navigated minimally invasive anterior approach and a conventional posterolateral approach for total hip arthroplasty: a

- randomized controlled trial. *J Orthop Res.* Feb 2013; 31(2): 288-94. PMID 22886805
16. Hsieh PH, Chang YH, Shih CH. Image-guided periacetabular osteotomy: computer-assisted navigation compared with the conventional technique: a randomized study of 36 patients followed for 2 years. *Acta Orthop.* Aug 2006; 77(4): 591-7. PMID 16929435
 17. Stiehler M, Goronzy J, Hartmann A, et al. The First SICOT Oral Presentation Award 2011: imageless computer-assisted femoral component positioning in hip resurfacing: a prospective randomised trial. *Int Orthop.* Apr 2013; 37(4): 569-81. PMID 23385606
 18. Xie C, Liu K, Xiao L, et al. Clinical Outcomes After Computer-assisted Versus Conventional Total Knee Arthroplasty. *Orthopedics.* May 2012; 35(5): e647-53. PMID 22588405
 19. Rebal BA, Babatunde OM, Lee JH, et al. Imageless computer navigation in total knee arthroplasty provides superior short term functional outcomes: a meta-analysis. *J Arthroplasty.* May 2014; 29(5): 938-44. PMID 24140274
 20. Blakeney WG, Khan RJ, Palmer JL. Functional outcomes following total knee arthroplasty: a randomised trial comparing computer-assisted surgery with conventional techniques. *Knee.* Mar 2014; 21(2): 364-8. PMID 24703685
 21. Lützner J, Dixel J, Kirschner S. No difference between computer-assisted and conventional total knee arthroplasty: five year results of a prospective randomised study. *Knee Surg Sports Traumatol Arthrosc.* Oct 2013; 21(10): 2241-7. PMID 23851969
 22. Beyer F, Pape A, Lützner C, et al. Similar outcomes in computer-assisted and conventional total knee arthroplasty: ten year results of a prospective randomized study. *BMC Musculoskelet Disord.* Aug 18 2021; 22(1): 707. PMID 34407776
 23. Cip J, Widemschek M, Luegmair M, et al. Conventional versus computer-assisted technique for total knee arthroplasty: a minimum of 5-year follow-up of 200 patients in a prospective randomized comparative trial. *J Arthroplasty.* Sep 2014; 29(9): 1795-802. PMID 24906519
 24. Hsu RW, Hsu WH, Shen WJ, et al. Comparison of computer-assisted navigation and conventional instrumentation for bilateral total knee arthroplasty: The outcomes at mid-term follow-up. *Medicine (Baltimore).* Nov 2019; 98(47): e18083. PMID 31764842
 25. Song EK, Agrawal PR, Kim SK, et al. A randomized controlled clinical and radiological trial about outcomes of navigation assisted TKA compared to conventional TKA: long-term follow-up. *Knee Surg Sports Traumatol Arthrosc.* Nov 2016; 24(11): 3381-3386. PMID 26831857
 26. Cip J, Obwegeser F, Benesch T, et al. Twelve-Year Follow-Up of Navigated Computer-Assisted Versus Conventional Total Knee Arthroplasty: A Prospective Randomized Comparative Trial. *J Arthroplasty.* May 2018; 33(5): 1404-1411. PMID 29426792
 27. Farhan-Alanie OM, Altell T, O'Donnell S, et al. No advantage with navigated versus conventional mechanically aligned total knee arthroplasty-10 year results of a randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc.* Mar 2023; 31(3): 751-759. PMID 36166095
 28. Kim YH, Park JW, Kim JS. Computer-navigated versus conventional total knee arthroplasty a prospective randomized trial. *J Bone Joint Surg Am.* Nov 21 2012; 94(22): 2017-24. PMID 23052635
 29. Hoppe S, Mainzer JD, Frauchiger L, et al. More accurate component alignment in navigated total knee arthroplasty has no clinical benefit at 5-year follow-up. *Acta Orthop.* Dec 2012; 83(6): 629-33. PMID 23140107
 30. Yaffe M, Chan P, Goyal N, et al. Computer-assisted versus manual TKA: no difference in clinical or functional outcomes at 5-year follow-up. *Orthopedics.* May 2013; 36(5): e627-32. PMID 23672916
 31. Hoffart HE, Langenstein E, Vasak N. A prospective study comparing the functional outcome of computer-assisted and conventional total knee replacement. *J Bone Joint Surg Br.* Feb 2012; 94(2): 194-9. PMID 22323685
 32. Dyrhovden GS, Fenstad AM, Furnes O, et al. Survivorship and relative risk of revision in computer-navigated versus conventional total knee replacement at 8-year follow-up. *Acta Orthop.* Dec 2016; 87(6): 592-599. PMID 27775460

33. Antonios JK, Kang HP, Robertson D, et al. Population-based Survivorship of Computer-navigated Versus Conventional Total Knee Arthroplasty. *J Am Acad Orthop Surg*. Oct 15 2020; 28(20): 857-864. PMID 31934926
34. Webb ML, Hutchison CE, Sloan M, et al. Reduced postoperative morbidity in computer-navigated total knee arthroplasty: A retrospective comparison of 225,123 cases. *Knee*. Jun 2021; 30: 148-156. PMID 33930702
35. Gelalis ID, Paschos NK, Pakos EE, et al. Accuracy of pedicle screw placement: a systematic review of prospective in vivo studies comparing free hand, fluoroscopy guidance and navigation techniques. *Eur Spine J*. Feb 2012; 21(2): 247-55. PMID 21901328
36. Shin BJ, James AR, Njoku IU, et al. Pedicle screw navigation: a systematic review and meta-analysis of perforation risk for computer-navigated versus freehand insertion. *J Neurosurg Spine*. Aug 2012; 17(2): 113-22. PMID 22724594
37. Staartjes VE, Klukowska AM, Schröder ML. Pedicle Screw Revision in Robot-Guided, Navigated, and Freehand Thoracolumbar Instrumentation: A Systematic Review and Meta-Analysis. *World Neurosurg*. Aug 2018; 116: 433-443.e8. PMID 29859354
38. Perdomo-Pantoja A, Ishida W, Zygorakis C, et al. Accuracy of Current Techniques for Placement of Pedicle Screws in the Spine: A Comprehensive Systematic Review and Meta-Analysis of 51,161 Screws. *World Neurosurg*. Jun 2019; 126: 664-678.e3. PMID 30880208
39. Laine T, Lund T, Ylikoski M, et al. Accuracy of pedicle screw insertion with and without computer assistance: a randomised controlled clinical study in 100 consecutive patients. *Eur Spine J*. Jun 2000; 9(3): 235-40. PMID 10905443
40. Rajasekaran S, Vidyadhara S, Ramesh P, et al. Randomized clinical study to compare the accuracy of navigated and non navigated thoracic pedicle screws in deformity correction surgeries. *Spine (Phila Pa 1976)*. Jan 15 2007; 32(2): E56-64. PMID 17224800
41. Villard J, Ryang YM, Demetriades AK, et al. Radiation exposure to the surgeon and the patient during posterior lumbar spinal instrumentation: a prospective randomized comparison of navigated versus non-navigated freehand techniques. *Spine (Phila Pa 1976)*. Jun 01 2014; 39(13): 1004-9. PMID 24732833
42. Arand M, Hartwig E, Kinzl L, et al. Spinal navigation in tumor surgery of the thoracic spine: first clinical results. *Clin Orthop Relat Res*. Jun 2002; (399): 211-8. PMID 12011712
43. Van Royen BJ, Baayen JC, Pijpers R, et al. Osteoid osteoma of the spine: a novel technique using combined computer assisted and gamma probe-guided high-speed intralesional drill excision. *Spine (Phila Pa 1976)*. Feb 01 2005; 30(3): 369-73. PMID 15682022
44. Smitherman SM, Tatsui CE, Rao G, et al. Image-guided multilevel vertebral osteotomies for en bloc resection of giant cell tumor of the thoracic spine: case report and description of operative technique. *Eur Spine J*. Jun 2010; 19(6): 1021-8. PMID 20069317
45. American Academy of Orthopaedic Surgeons. Surgical management of osteoarthritis of the knee. 2022. <https://www.orthoguidelines.org/topic?id=1041>. Accessed February 13, 2024.

Review/Revision/Approval History	
Date	Description
07/01/2024	New Policy
04/27/2026	Reviewed and Approved by Policy Committee

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult appropriate healthcare providers for medical advice, care, and treatment. Benefits

and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's benefit plan, effective when services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Including or excluding a procedure, diagnosis, or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as they apply to an individual member.

Mountain Health Co-Op makes no representations and accepts no liability regarding the content of any external information cited or relied upon in this policy. Mountain Health Co-Op updates its Coverage Policies regularly and reserve

es the right to amend these policies and give notice per State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Mountain Health Co-Op.

"Mountain Health Co-Op" and its accompanying logo and marks are protected and registered trademarks of Mountain Health Co-Op. The content of this Service is proprietary and protected by copyright. You may access the copyrighted content of this Service only for purposes outlined in these Conditions of Use.

© CPT Only – American Medical Association